THE BRYOLOGIST

Quarterly Journal of the American Bryological and Lichenological Society

Editor
DALE M. J. MUELLER

Volume 79
1976

Table of Contents

ALBERTINE, K. H., N. C. MARAVOLO & H. KAUSTINEN. Effects of gibberellin and several growth retardants on plastid ultrastructure in the hepatic *Marchantia polymorpha* 22

ANDERSON, L. E. Obituary—Antero Vaarama, 1912–1975 377

ANDRUS, R. E. & E. F. LAYSER. *Sphagnum* in the northern Rocky Mountains of the United States 508

BADHE, P. D. Free amino acids in two species of Anaptychia (lichens) 354

BAYFIELD, N. G. Effects of substrate type and microtopography on establishment of a mixture of bryophytes from vegetative fragments 199

BOWLER, P. A. & C. W. SMITH. *Ramalina farinacea* in Hawaii 499

BRASSARD, G. R. The mosses of Northern Essesmere Island, Arctic Canada. III. New or additional records 480

———. *Seligeria brevifolia*, a moss new to North America 106

———. Review 547

———, D. P. WEBER & A. J. FIFE. New or additional moss records from Newfoundland II 84

BREIL, D. A. & S. M. MOYLE. Bryophytes used in construction of bird nests 95

BRODO, I. M. *Lichenes canadenses exsiccati*: Fascicle II 385

———. A new combination for *Cladonia stellaris* 363

BUCK, W. R. & H. CRUM. Revision of the genus *Rozea* (Musci) 406

CHURCHILL, S. P. Contributions toward a moss flora of Nebraska 241

CROSBY, M. R. The correct name for the moss *Duseniella* 246

———. Recent literature on mosses—104–105 ..... 107, 535

———. The relationship of the mosses *Hookeria wrightii* and *Stenodictyon sericeum* 83

CULBERSON, W. L. & C. F. CULBERSON. Obituary—A tribute to Yasuhiko Asahina 258

———. Recent literature on lichens—93–94 ..... 113, 523

CULBERSON, C. F. & T. L. ESSLINGER. 4-O-Methylolivetoric and loxodellic acids: New depsides from new species of brown parmeliae 42

DELGADILLO, C. M. *Indusiella andersonii* sp. nov. in Northern Africa 98

——— & W. D. REESE. Notes on tropical mosses of Mexico 511

DOUGLAS, G. W. & D. H. VITT. Moss-lichen flora of St. Elias-Kluane Ranges, southwestern Yukon 437

EGAN, R. S. New and additional lichen records from Texas 224

———. Review 549

ENGEL, J. J. Austral Hepaticae VI. Some new species and new combination of taxa from southern Chile 514

FRENCH, J. C. & D. J. PAOLILLO, JR. Effect of light and other factors on capsule expansion in *Funaria hygrometrica* 457

FULFORD, M. Recent literature on hepatics—73–75 ..... 253, 370, 531

GALLOWAY, D. J. Obituary—William Martin, 1886–1975 374

GODFREY, J. D. *Schofieldia*, a new hepatic from the Pacific Northwest 314

GROSSMAN, H. H. & C. J. HILLSON. The effect of chronic low-level gamma radiation on *Polytrichum commune*. II. Regeneration in leaf and stem-leaf cuttings 488

HALE, M. E., JR. *Hypotrachyna showmanii*, a new lichen from eastern North America 78
HERMANN, F. J. Additions to the bryophyte flora of Colorado 215
———. Recopilacion de los musgos de Bolivia 125
HICKS, M. L. *Lophozia heterocolpa*, new disjunct for North Carolina 222
HONG, W. S. Annotated checklist of the hepatics of Idaho 422
——— & D. H. VITT. A list of hepaticae from west-central Alberta 180
HORTON, D. G. & B. M. MURRAY. *Encalypta brevipes* and *E. mutica*, *gymnostomous* species new to North America 321
IRELAND, R. R. Alar cells of *Rhacomitrium lawtonae* 238
——— & H. CRUM. Distribution of *Grimmia cribrosa* in North America 355
JOHNSEN, I. & U. SØCHTING. Distribution of cryptogamic epiphytes in a Danish city in relation to air pollution and bark properties 86
KARUNEN, P., N. VALANNE & R. E. WILKINSON. Influence of S-Ethyl dipropylthiocarbamate on growth, chlorophyll and carotenoid production and chloroplast ultrastructure of germinating *Polytrichum commune* spores 332
KROG, H. & T. D. V. SWINSCOW. *Pyxine azorea* Nyl. is *Parmelia caroliniana* Nyl. 506
LANGE, B. *Sphagnum pylaesii* in Greenland 47
LEBLANC, F. Review 264
MAGILL, R. E. Mosses of Big Bend National Park, Texas 269
——— & R. KIRKPATRICK. Notes on Kansas mosses 101
MCFARLAND, K. D. & W. A. WISTENDAHL. Relationship of terrestrial mosses to forested flood plains in southeastern Ohio 64
MILLER, H. A. Review 548
MILLER, N. G. & L. J. H. AMBROSE. Growth in culture of wind-blown bryophyte gametophyte fragments from Arctic Canada 55
MUELLER, D. M. J. Review 264
NAIR, H. & V. RAGHAVAN. Cytokinin induced bud formation in *Microdus miquelianus* 495
NASH, T. H. III. Sensitivity of lichens to nitrogen dioxide fumigations 103
OATHAGEN, H. Nomenclatural note on *Cladonia nemoxyna* 242
OSORIO, H. S. Contribution to the lichen flora of Argentina. VIII. Lichens from Punta Lara, Buenos Aires Province 358
PAOLILLO, D. J., JR. & M. CUKIERSKI. Wall developments and coordinated cytoplasmic changes in spermatogenous cells of *Polytrichum* (Musci) 466
PURSELL, R. A. Index to Volume 79 552
———. On the typification of certain taxa and structural variation within the *Fissidens bryoides* complex in eastern North America 35
PYATT, F. B. Lichen ecology of metal spoil tips: Effects of metal ions on ascospore viability 172
RASTORFER, J. R. Caloric values of three Alaskan-Arctic mosses 76
———. Review 122
REDFEARN, P. L., JR. A *Trichostomum* from Texas new to the United States 80
REESE, W. D. & D. GRIFFIN, III. *Syrrhopodon annotinus* (Musci; Calymperaceae), a new species from Amazonas, Brazil 518
RUNDEL, P. W. & P. A. BOWLER. *Ramalina leptocarpha* and *R. subleptocarpha*: A fertile-sorediate species pair 364

SAADAWI, W. S. Some mosses from Kuwait 515
SCHUSTER, R. M. Review 380
SHEARD, J. W. Review 382
SHERIDAN, R. P., C. SANDERSON & R. KERR. Effects of pulp mill emissions on lichens in the Missoula Valley, Montana 248
SHOWMANN, R. E. Seasonal growth of *Parmelia caperata* 360
SMITH, G. L. Neotropical Polytrichaceae IV 93
———. Neotropical Polytrichaceae V 521
STEERE, W. C. Identity of *Pterygoneurum arcticum* with *P. lamellatum* 221
——— & G. R. BRASSARD. *Schistidium holmenianum*, sp. nov. from Arctic North America 208
STOTLER, R. E. Saxicolous Bryophyte and macrolichen associations in southern Illinois. I. Little Grand Canyon, Jackson County 1
SWANSON, E. S., W. H. ANDERSON, J. L. GELLERMAN & H. SCHLENK. Ultrastructure and lipid composition of mosses 339
THOMSON, J. W. & T. H. NASH III. Three new lichens from the southwest: *Xanthoria concinna* sp. nov., *Lecanora collatolica* sp. nov. and *Toninia conglomerata* 350
THOMAS, P. E. & P. L. REDFEARN, JR. Bryophytes of the interior highlands XVI. New distributional records for Arkansas and Missouri 100
UDAR, P. & D. K. SINGH. A new *Cyathodium* from India 234
VALANNE, N., S. TOIVONEN & R. SAARINEN. Ultrastructural changes in germinating *Dicranum scoparium*: A moss containing protein storage material 188
VITT, D. H. Mosses new to the Yukon from the Ogilvie Mountains 501
——— & W. B. SCHOFIELD. *Seligeria careyana*, a new species from the Queen Charlotte Islands, Western Canada 231
WETMORE, C. M. Macrolichens of Big Bend National Park, Texas 296
ZANDER, R. H. & E. HEGEWALD. *Leptodontiella*, gen. nov. and *Leptodontium* from Peru 16
ZANDER, R. H. Notes on Pottiaceae in Middle America 227

Russian chapter of the Sullivant Moss Society in 1926.

* * * *

Varieties of the Philonotis *species in Finland.*—In a selection of herbarium material from the University of Helsinki sent by Dr. T. J. Koponen, I have found the following *Philonotis* varieties: *P. fontana* (Hedw.) Brid. var. *adpressa* (Ferg.) Limp. and var. *tomentella* (Mol.) Dix.; *P. caespitosa* Wils. ex Jur. var. *adpressa* Dism.* and var. *laxa* Lske.& Warnst.*; *P. calcarea* (B.&S.) Schp. var. *laxa* Dism. and var. *orthophylla* Schiffn.*; and *P. seriata* Mitt. var. *adpressa* Lske.&Monk.* and var. *laxa* J.H.Field.*

I am informed by P. Isoviita that *Philonotis calcarea* var. *laxa* was reported from Finland by H. Rancken in 1914.

The taxa marked with an asterisk are likely to be new records for the Finnish bryoflora. —J. H. Field, *North Warwickshire College of Technology and Art, England.*

* * * *

Summer Course in Bryology.—The Lake Itasca Forestry and Biological Station, University of Minnesota, will offer a course in bryology, 17 July–20 August. The course will be taught by Frank D. Bowers, Biology Department, University of Wisconsin-Stevens Point, Stevens Point, Wisconsin. For further information contact Dr. D. F. Parmelee, Field Biology Program, University of Minnesota, Minneapolis, Minnesota 55455.

* * * *

A 1973–4 collection of mosses from western Australia, Tasmania, Queensland and Fiji by D. H. Norris is now fully curated and available to monographers. This collection consists of 21,000 numbers and, especially for Tasmania and Queensland, seems to represent a nearly complete set of the known flora, as well as many additions. Loan requests should be made to Daniel H. Norris, Department of Botany, Humboldt State University, Arcata, California 95521.

* * * *

Statement of ownership, management and circulation of The Bryologist required by Act of 12 August 1970: Section 3685, Title 39, United States Code, filed 30 September 1976.

The Bryologist is published quarterly at 2345 Tower Grove Avenue, St. Louis, Missouri 63110. General Business Offices: Missouri Botanical Garden, 2345 Tower Grove Avenue, St. Louis, Missouri 63110. Publisher: American Bryological & Lichenological Society, Inc. Editor: D. M. J. Mueller, Dept. of Biology, Texas A&M University, College Station, Texas 77843. Owner: American Bryological & Lichenological Society, Inc., 2345 Tower Grove Avenue, St. Louis, Missouri 63110.

The known bondholders, mortgagees and other security holders owning or holding 1% or more of the total amount of bonds, mortgages or other securities are: none.

Purpose, function and non-profit status of this organization and the exempt status for Federal income tax purposes have not changed during the preceding 12 months.

Circulation: Total number of copies printed: average number of copies each issue during preceding 12 months, 1150; number of copies of single issue published nearest to filing date, 1150. Sales through dealers and carriers, street vendors and counter sales: none. Paid mail subscriptions: average number of copies each issue during preceding 12 months, 961; actual number of copies of single issue published nearest to filing date, 979. Free distribution: 9. Total distribution: average number of copies at each issue during preceding 12 months, 970; actual number of copies of single issue published nearest to filing date, 988. Copies not distributed: average number of copies each issue during preceding 12 months, 180: actual number of copies of single issue published nearest to filing date, 162.

I certify that the statements made by me above are correct and complete.

Signed,
Marshall R. Crosby, Business Manager.

Volume 79, Number 3 (pages 269–384) was issued 15 October 1976

Index to Volume 79 of The Bryologist[1, 2]

Acacia spp., 269-
Acanthocladium subnitidum, 157
Acarospora chlorophana, 447; glaucocarpa, 447; oxytona, 448; veronensis, 448
Acaulon schimperianum, 281
Acer grandidentatum, 269-; negundo, 64-; pseudoplatanus, 86-, 172-; saccharinum, 64-
Acidodontium brachypodium, 139, 162; exaltatum, 139; lonchotrachylon, 139, 162; longifolium, 139, 162; macropoma, 139; pallidum, 139; ramicola, 139; spinicuspis, 139
Acrocryphaea gardneri, 148; julacea, 148
Acroporium pungens, 157, 168
Adelothecium bogotense, 151
Africa. Indusiella andersonii in northern, 98
Agave americana, 269-; lechuguilla, 269-
Agrestia cyphellata, 386; hispida, 386, 387*
Agropyron yukonense, 437-
air pollution. distribution of cryptogamic epiphytes in a Danish city in relation to, 86; effects of pulp mill emission on lichens, 248; lichen sensitivity to nitrogen dioxide, 103
Alaska. caloric values of three Alaska-Arctic mosses, 76; distribution of Encalypta brevipes and E. mutica in, 321, 328*; distribution of Grimmia cribosa in, 356, 356*; Pterygoneurum lamellatum in, 221; Schistidium holmenianum in, 208
Alberta. *See* Canada
ALBERTINE, K. H., N. C. MARAVOLO & H. KAUSTINEN. Effects of Gibberellin and Several Growth Retardants on Plastid Ultrastructure in the Hepatic Marchantia polymorpha, 22
Alectoria abbreviata, 392; capillaris, 386; fremontii, 248; lanestris, 448; nadvornikiana, 387; nidulifera, 387; nigricans, 448; ochroleuca, 448; oregana, 392; pubescens, 448; tenuis, 387
alectorialic acid, 385-
alectoronic acid, 224-
ALLISON, K. W. & J. CHILD. The Liverworts of New Zealand (review), 548
Alnus crispa var. lacinata, 437-
Aloina brevirostris, 440, 481; hamulus, 281; rigida, 216, 281, 516; sp., 55-; spp., 98-
Aloinella boliviana, 132
Amblyodon dealbatus, 440
Amblystegiaceae, 154, 218, 292
Amblystegium loeskei, *155*, *161*; riparium, 1-; rotae, 219; serpens, 1-, 86-, 95-, 292, 440; varium, 1-, 95-, 154
AMBROSE, L. J. H. *See* MILLER, N. G. & L. J. H. AMBROSE
Amelanchier arborea, 222-
amino acids in two species of Anaptychia, 354
Amphidium cyathicarpum, 145; cyathicarpum var. fragilifolium, 145; lapponicum, 321-, 482
Anacamptodon cubensis, 153
Anacolia laevisphaera, 142, 289; setifolia, *144*, *161*
Anaptychia diademata, 354, 358; diademata fo. brachyloba, 358; domingensis, 358; neoleucomelaena, 103-; obscurata, 354-; podocarpa, 354
Anastrophyllum assimile, 425; donianum, 425; hellerianum, 183, 425; michauxii, 183, 425; minutum, 183, 425, 434
Ancistrodes **genuflexa**, comb. nov., 247
ANDERSON, L. E. ANTERO VAARAMA (1912–1975), 377
ANDERSON, W. H. *See* SWANSON, E. S. et al.
Andreaea angustifolia, 127; arachnoidea, 127; arachnoidea var. gracilis, 127; barbuloides, 127; clavata, 127; clavata var. tenuior,

[1] Compiled by R. A. Pursell, Department of Biology, Buckhout Laboratory, The Pennsylvania State University, University Park, PA 16802.

[2] New names and combinations are in **boldface** type. An asterisk (*) indicates an illustration is on the page cited. A hyphen (-) following the page citation indicates that the entry is mentioned incidentally or is mentioned more than once in the article beginning on the page cited. A page number in *italics* indicates that the taxon signified is given as a synonym on the page cited. Names of authors of contributions and of publications reviewed are in LARGE and SMALL CAPITALS.

127; dissitifolia, 127; erythrodictyon, 127; laticuspis, 127; lorentziana, 127; nivalis, 502; obovata, 502; pseudosubulata, *127*, *161*; robusta, 127; rupestris, 440, 482; rupestris var. acuminata, 502; rupestris var. rupestris, 502; squarrosa, 127; striata, 127; subenervis, 127; subulata, 127, 161; tunariensis, 127; turgescens, 127; vilocensis, 127
Andreaeaceae, 127
ANDRUS, R. E. & E. F. LAYSER. Sphagnum in the northern Rocky Mountains of the United States, 508
Aneura pinguis, 186, 432
Aneuraceae, 186, 432
Anisothecium campylophyllum, 128, 163; hookeri, 128, 163; laxirete, 128, 163; macrostomum, 128, 161, 163; spiripes, 163; submacrostomum, 128, 163
Anoectangium aestivum, 133, 161, 227, 228*; brachyphyllum, 133; compactum, *133*, *161*; euchloron, 133, 227-; herzogii, 133; lechlerianum, 133; lineare, 133, 161, 169; mandonianum, *133*, *161*; obtusifolium, 281; pflanzii, 133; tenuinerve, 482
Anomobryum filiforme, 139; humillimum, 139, 162; obtusatissimum, 139, 162; orbiculatum, 139; prostratum, *139*, *161*; robustum, 139; semiovatum, 139; soquense, 139, 161, 162
Anomodon attenuatus, 1-, 95-, 241; fragillimus, 154; minor, 95-; rostratus, 1-, 241
Anthelia juratzkana, 185
Antheliaceae, 185
antheridia of Polytrichum, 468*, 470*, 472*, 474*, 476*, 478*
Anthracothecium goniostomum, 358
Anthriscus sylvestris, 172-
Aongstroemia julacea, 129; macrostoma, *128*, *161*; nanocarpa, *131*, *161*
Aptychella caudata, 157; linearifolia, 157; prolifera, *157*, *161*; proligera, 157, 167; proligera var. chlorophyllosa, 157
Aptychus brachyacrus, *158*, *161*; chlorocormus, *158*, *161*
Aquilegia canadensis, 222-
Arbutus xalapensis, 269-
Arctic America. *See* Alaska, Canada and Greenland
Arctoa fulvella var. andersonii, 502
Arctostaphylos, 437-
Argentina. lichens from Punta Lara, Buenos Aires Province, 358
Aristida spp., 269-
Arizona. distribution of Grimmia cribosa in, 356, 356*; Pterygoneurum lamellatum in, 221; sensitivity of lichens [collected in Arizona] to nitrogen dioxide, 103-; Toninia conglomerata reported from, 353
Arkansas. new distributional records of bryophytes for, 100
Arnellia fennica, 185
Artemisia frigida, 437-
Arthopyrenia punctiformis, 388
Arthrorhaphis citrinella, 448
Arundo donax, 269-
ARZENI, C. B. & B. N. MCKNIGHT. Buxbaumia aphylla, A New Report for Illinois, 266
Asahina, Y., 258*
ascospore viability, 172
Aspicilia calcarea, 386
Asplenium montanum, 222-; rhizophyllum, 1-; trichomanes, 222-
associations of bryophytes and macrolichens, 1
Asterella lindenbergiana, 432; ludwigii, 432; saccata, 432; tenella, 1-
Astomiopsis amblyocalyx, 128
Astomum drummondii, *384*; muhlenbergianum, 281; phascoides, *384*; × phascoides, 281
Atractylocarpus longisetus, 129, 165; **spiripes**, comb. nov., 129, 165
atranorin, 78-, 224-, 296-, 353, 385-, 506
Atrichum angustatum, 1-, 266-; elamellosum, 160, 162; **nigricans**, comb. nov., 160, 162; polycarpum, 160, 162; undulatum, 95-
Aulacomniaceae, 142
Aulacomnium acuminatum, 440; heterostichum, 1-; marginatum var. andinum, *142*, *161*; palustre, 95-, 101, 199-, 201*, 241, 440; palustre var. andinum, 142, 161; turgidum, 440, 482
Australia. Parmelia loxodella in Tasmania, 43

Bacidia alutacea, 358; obscurata, 448; sphaeroides, 448
BADHE, P. D. Free Amino Acids in Two Species of Anaptychia (Lichens), 354
Baeomyces radiatus β. nemoxynus, *245*; roseus, 388; rufus, 172-
barbatic acid, 385-
barbatolic acid, 385-
Barbella tenuissima, 149, 164
Barbula, 98-; acuta, 215-, 440; acuta var. bescherellei, 281; aenea, *229*; amblyacra, *133*, *161*; andicola, *136*, *161*; apiculata, *133*, *161*; apiculata fo. laxa, *133*; apiculata var. breviseta, *133*, *161*; apiculata var. gracilis, *133*, *161*; austro-revoluta, 133; bescherellei, 216; boliviana, 133, 164; boliviensis, *133*, *161*; brachyphylla, 216; brunnea, *136*, *161*; campylocarpa, 133, 169; chocayensis, 133; decolorans, 133, 161, 163; fallax, 101, 282, 440; flexifolia, *133*, *161*; fusca, 133, *133*, *161*; germainii, *135*, *161*; glaucescens, 133; glau-

cescens var. latifolia, 133; humboldtii, 133; icmadophila, 440, 482; imperfecta, 133; inaequalifolia, 229; jamesonii, *133*, *161*; johansenii, *483*; laevigata, 133; longirostris, 133, 169; lurida, 133, 161; mandonii, 133, 161; mniadelphus, *136*, *161*; pachygastrella, *135*, *161*; paludicola, *136*, *161*; perexilis, *134*, *161*; pflanzii, 133; pflanzii var. falcatula, 133; polyseta, *136*, *161*; punae, 133; pygmaeola, *135*, *161*; rectifolia, 133; replicata, 133, 161; rothii, 133, 161; sp., 133, 164, 516; spiralis, 282; subandreaeoides 215-; subglaucescens, 133; tristis, 133, 161; unguiculata, 133, 216; vinealis, 516; viridula, *136*, *161*

bark properties. distribution of cryptogamic epiphytes in relation to air pollution and, 86

Bartramia, 122-; ambigua, 142; ambigua var. penicillatula, 142; asperrima, *142*, *161*; auricola, 142; brachyphylla, *144*, *161*; breviseta, *143*, *161*; brittoniae, *143*, *161*; brotheri, 142; cacaltayae, 142, 161, 162; defolians, 142; didymocarpa, 142; filiramea, *144*, *161*; flavicans, 142; fragilifolia, 142, 161; fragilifolia fo. compacta, 142; fragilifolia fo. latifolia, 142; glauca, *142*, *161*; guyabayana, *144*, *161*; hampeana, 142, 161; inflata, *142*, *161*; ithyphylla, 85, 321-, 440, 482, 502; ithyphylloides, 143; jamesonii, *144*, *161*; macrocarpa, *143*, *162*; macropoma, 143; mathewsii, 143; micrangia, 143; minutissima, *144*, *162*; mniocarpa, *143*, *162*; mollis, 143; perpumila, 143; perpusilla, 143; pilicuspis, 143; pinnulata, *144*, *162*; polytrichoides, 143; pomiformis, 222-, 441; potosica, 143; pruinata, 143; pugionifolia, *144*, *162*; rigidissima, 143; rosea, 143; scorpioides, *143*, *162*; secunda, 143; secundifolia, *143*, *162*; squarrosa, 143; subithyphylla, 143; thrausta, 143; tomentosa, *143*, *162*; wedellii, *142*, *162*; wedellii fo. irrorata, *142*, *162*; xanthochaeta, 143

Bartramiaceae, 142, 218, 289

Bayfield, N. G. Effects of Substrate Type and Microtopography on Establishment of a Mixture of Bryophytes from Vegetative Fragments, 199

Bazzania ambigua, 425, 435; denudata, 425; tricrenata, 266-, 425; trilobata, 1-, 425

Bellis perennis, 172-

Betula, 241-; glandulosa, 437-; nigra, 1-; spp., 172-

Bhutan. distribution of Rozea in, 417

Biatorina baeomma, *390*

Big Bend National Park, Texas, 270*, 274*, 298*; key to macrolichens in, 300; keys to mosses in, 269

bird nests. bryophytes used in, 95

Blasiaceae, 186, 431

Blasia pusilla, 186, 431

Blechnum spp., 172-

Blepharostoma arachnoideum, 424; trichophyllum, 182, 424; trichophyllum subsp. brevirete, 182, 424

Blepharostomaceae, 182, 424

Blindia, 231-; acuta, 441, 482

Bolivia. notes on Pottiaceae in, 227; recopilación de los musgos de, 125

boninic acid, 350

Bouteloua romosa, 269-

Bowler, P. A. & C. W. Smith. Ramalina farinacea in Hawaii, 499

Bowler, P. A. *See* Rundel, P. W. & P. A. Bowler

Brachydontium, 231-

Brachymenium acuminatum, 139; barbuloides, 139, 162; dimorphum, 139; flexipilum, 139; jamesonii, *139*, *162*; mexicanum, 288; radiculosum, 139; speciosum, 139, 162; systylium, 288

Brachymitrion jamesonii, *138*, *162*

Brachytheciaceae, 155, 219, 293

Brachythecium, 406-; acuminatum, 95-; albicans, 441; asperrinum, 219; bolivioplumosum, 155; calcareum, 441; calliergonoides, 155; cavifolium, 155; chocayae, 155; cochleare, 155; collinum, 293, 441; erythrorrhizon, 219; fendleri, 293; fissidentoides, 155; flaccum, 155; frigidum, 441; grandirete, 155; groenlandicum, 482, 502; herzogii, 155, 162; hirtipes, 155, 167; julaceum, 155; leibergii, 219, 502; lescuraeoides, 155; longisetum, 155; mollerameum var. delicatulum, 155; oxycladon, 64-, 95-; plumosum, 219; praelongum, 155; riparium, 155; rivulare, 64-; rivulariforme, 155; rutabulum, 95-, 155, 219; salebrosum, 1-, 95-, 219, 441; scaberrimum, *155*, *162*; scabripes, *156*, *162*; serrulatum, 1-; starkei, 220; stereopoma, 155, 409; subjulaceum, 155; sulphureum, 155; tenuipinnatum, 155; trachypodioides, 155; turgidum, 441, 482, 502; velutinum var. venustum, 220

Brassard, G. R. (review) M. Kuc, Bryogeography of Expedition Area, Axel Heiberg Island, N.W.T., Canada, 547

Brassard, G. R. Seligeria brevifolia, a Moss New to North America, 106

Brassard, G. R. The Mosses of Northern Ellesmere Island, Arctic Canada. III. New or Additional Records, 480

Brassard, G. R., D. P. Weber & A. J. Fife.

New or Additional Moss Records from Newfoundland II, 84
BRASSARD, G. R. *See also* STEERE, W. C. & G. R. BRASSARD
Braunia argyrotricha, *147*, *162*; canescens, *147*, *162*; cirrhifolia, 147, 162; crenulata, *162*; divaricatula, 147; laxifolia, 147; plicata, 147; secunda, 147, 291; subplicata, 147
Brazil. neotropical Polytrichaceae in, 521
Breidleria amabilis, *159*
BREIL, D. A. & S. M. MOYLE. Bryophytes Used in Construction of Bird Nests, 95
Breutelia, 122-; anacolioides, 143; auricolor, 143; boliviensis, 143; brevifolia, 143; breviseta, 143, 161, 166; brittoniae, 143, 161; bryocarpa, 143, 162; crispula, *143*, *162*; gertrudis, 143; harpophylla, 143; hasskarliana, 143; inclinata, 143; integrifolia, 143; lorentzii, 143, 162; macrocarpa, 143, 162; macrotheca, 143; minuta, 143; mniocarpa, 162; nigrescens, 143; nutans, *143*, *162*; patens, 143; scorpioides, 143, 162; secundifolia, 143, 162; straminea, 143; subdisticha, 143; tomentosa, 143, 162, 166; undulata, 143; wainioi, 143
British Columbia. *See* Canada
BRODO, I. M. A New Combination for Cladonia stellaris, 363
BRODO, I. M. Lichenes Canadenses Exsiccati: Fascicle II, 385
Brotherella tenuirostris, 95-
Bruchia texana, 241
Bryaceae, 139, 217, 288
Bryhnia graminicolor, 95-; novae-angliae, 95-; pflanzii, 155
Bryoandersonia illecebra, 1-, 266-
Bryobrittonia pellucida, 441
Bryodusenia, *247*; genuflexa, *247*
Bryoerythrophyllum campylocarpum, 133, 163, 169; jamesonii, 133, 161, 163, 164; recurvirostre, 229-, 282, 441; recurvirostre var. **aeneum**, stat. nov., 229
bryophyte ecology. *See* ecology
Bryosedgwickia, 406-
Bryum albicans, *141*, *162*; albidum, 139; algovicum, 441; angustifolium, *139*, *162*; apophysatum, 139; archangelicum, 502; arcticum, 482; argenteum, 55-, 86-, 139, 162, 441, 482; fo. ochrotheca, 139, 162, var. argenteum, 288; argenteum var. lanatum, 139, 162, 288; atenense, 139; barbuloides, *139*, *162*; bediloma, 139; bicolor, 517; billardierei, 289; bimum, *140*, *162*, 482; bolivianum, 139, 162; brachypodium, *139*, *162*; caespiticium, 289, 441; calophyllum, 482; candicans, 139; capillare, 86-, 289; capillipes, *139*, *162*; capillipes var. dimorphum, *139*, *162*; capillipes var. subulatum, *139*, *162*; caulifolium, 139, 167; cavum, 139; challaense, 139; chryseum, 139; coloratum, 139; concavum, 139; congestum, 139; coronatum, 139; creberrimum, 217, 289, 441; cryophilum, 441, 482; cymbifolium, *139*, *162*; densifolium, 139; flexisetum, 140; fragilicuspis, 140; gemmiparum, 217; genucaule, 140; grandifolium, *142*, *162*; hamipilum, 140; humillimum, *139*, *162*; incacorralis, 140; laevigatum, 140; lapponicum, 84-; lindigianum, 140, 167; linearifolium, 140; lonchotrachylon, *139*, *162*; longedecurrens, 140; longifolium, *139*, *162*; malacophyllum, 140; mandonii, *140*, *162*; maynense, 140; micro-comosum, 140; micro-comosum fo. propagulifera, 140; nanophyllum, 140; neodamense, 482; nigro-purpureum, *140*, *162*; nitidulum, 482; niveum, 140; obtusatissimum, *139*, *162*; oediloma, 140; pallescens, 502; perserratum, 140; philonoteum, 140; proliferum, 140, 162; pseudotriquetrum, 95-, 140, 162, 441; pulchirete, 140; rupicola, 140; rusbyanum, *141*, *162*; salinum, 85; schisticola, *141*, *162*; sericeum, 140; soboliferum, 140; sp. 55-, 199-, 515; spininervium, 140; stenotrichum, 289, 482; stephanii, 140; subgenucaule, 140; subnanophyllum, 140; subsericeum, 140; terebellum, 140; turbinatum, 289; verrucosum, *142*, *162*; wrightii, 321-, 482, 502
BUCK, W. R. & H. CRUM. Revision of the Genus Rozea (Musci), 406
Buellia atrata, 389; dialyta, 389; epigaea, 389, 448; insignis, 448; moriopsis, *389*; papillata, 448; stellulata, 448; triphragmioides, 448; zahlbruckneri, 448
Burma. distribution of Rozea in, 420
Buxbaumia aphylla, 266, 321-

Calicium curtisii, *402*; populneum, *402*; schaereri, *389*; subquercinum, 389; trabinellum, 389; xylonellum *β*. C. trabinellum, *389*
California. distribution of Ramalina leptocarpha and R. subleptocarpha in, 366*, 367*, 368, 369; new species of lichen from, 352
Callicladium haldanianum, 95-
Callicostella integrifolia, 151, 164; microcarpa, 151; mosenii, 151; pallida, 151; rivularis, 151; scabripes, 151, 164; scabriuscula, 151; strumulosa, 151; subdepressa, 151
Calliergon cordifolium, 218, 441; giganteum, 441; luipichense, 154; richardsonii, 441; sarmentosum, 441, 482; stramineum, 47-, 441, 482; trifarium, 482

Calluna, 199-; vulgaris, 172-
Caloplaca americana, 358; cinnamomea, 448; cirrochroa, 448; fraudans, 448; holocarpa, 358, 448; jungermanniae, 448; marina, 389; sinapisperma, 390; stillicidiorum, 448; tetraspora, 448; xanthaspis, 358
caloric values of three Alaskan-Arctic mosses, 76
Calymperaceae, 132, 518
Calymperes bolivianum, *132*; emersum, 512; erosum, 512; levyanum, 512; lonchophyllum, 512; nicaraguense, 132, 512; richardii, 518
Calypogeia fissa, 425; integristipula, 425; muelleriana, 1-, 425; neesiana, 425; sp., 266-; sphagnicola, 425; suecica, 183, 425; trichomanes, 314-, 425
Calypogeiaceae, 183, 425
Calyptogon angustifolius, *136*, *162*
Calyptothecium duplicatum, 150; duplicatum var. integerrimum, 150
Campanula divaricata, 222-
Camptodontium fallax, 129, 163; flexipes, 129, 163
Campylium arcticum, 482; chrysophyllum, 95-, 293, 441; halleri, 218, 502; hispidulum, 95-, 154, 293; polygamum, 95-, 218, 441; polygamum var. latifolium, 154; radicale, 95-; stellatum, 76, 441
Campylodontium bolivianum, 156; onustum, *157*, *162*; sulcatum, *129*
Campylopodium krauseanus, *162*; sulcatum, *162*
Campylopus albidovirens, 129; alopecurus, 129; alopecurus var. bolivianus, 129; amboroensis, 129; andicola, 129; areodictyon, *129*, *162*; benedictii, 129; buchtienii, 129; cavifolius, 129; chrismarii, 129, 162; chrysodictyon, 129; clavatus, 129, 162; concolor, 129, 130, 162; cucullatifolius, 129; densicoma, 129, 163; filicuspis, 129; filifolius, 129, 162; flexuosus, 129, 162, 279; fulvus, 129; gertrudis, 129; harpophyllus, 129; heterophyllus, 129; humilis, *129*, *162*; incacorallis, 129; ingeniensis, 129; insignis, 129, 130; introflexus, 129, 162; jamesonii, 129; jugorum, 129; jugorum var. patulifolius, 129; julaceus, 129; lamellatus, *129*, *162*; lamellinervis, 129, 162; latinervis, *130*, *162*; laxiretis, 129; leptodus, *129*, *162*; leucognodes, 129, 162, 163; leucognodes var. gracilescens, 129; liliputanus, 129, 166; longisubulatus, 129, 162; longisubulatus var. squarrosus, 129; malagensis, 129; mandonii, 130; microtheca, 130; multicapsularis, 130, 163; nanofilifolius, 130, 163; occultus, 130; pelichucensis, 130; penicillatus, *129*, *162*; perexilis, 130, 163; perreduncus, 130, 163; porphyreodictyon, *129*, *162*; pseudodicranum, 130; ptychotheca, 130, var. circinatus, 130; renneri, 130, 162, var. latelimbata, 130; retroflexus, 130; richardii, 130, 166; rosulatus, 130; spirifolius, 130; spurio-concolor, 130, 163; subannotinus, 130; subcubitus, 130; subfimbriatus, 130; subfulvus, 130; subgriseus, 130; subincacorralis, 130; subjugorum, 130; subjugorum var. edithae, 130; subleucogaster, *129*, *162*; tablasensis, 130; thysanomitrioides, 130; totarae, 130; trichophorus, 130; trivialis, 130, 163; trollii, 130; tunariensis, 130; verticillatus, 130; yungarum, 130; zygodonticarpus, *129*, *162*;
Canada. distribution of Encalypta brevipes and E. mutica in Alberta and Yukon, 321, 328*; distribution of Grimmia cribosa in 356, 356*; distribution of Ramalina subleptocarpha in British Columbia, 367*, 368; growth in culture of wind-blown bryophyte gametophyte fragments from Arctic, 55; Hepaticae from West-Central Alberta, 180; Lichenes Canadenses Exsiccati [British Columbia, Manitoba, Newfoundland, Northwest Territories, Nova Scotia, Ontario, Quebec, Saskatchewan], 385; mosses new to the Yukon from the Ogilvie Mountains, 501; moss-lichen flora of S. Elias-Kluane Ranges, southwestern Yukon, 437; new or additional moss records from Newfoundland, 84; new or additional moss records from northern Ellesmere Island, 480; Schistidium holmenianum in, 212; Schofieldia in British Columbia, 318; Seligeria brevifolia, a moss new to North America [Newfoundland], 106; Seligeria careyana in the Queen Charlotte Islands, 232
Candelaria concolor, 302; fibrosa, 302; sp., 302
Candelariella aurella, 448; terrigena, 448; vitellina, 448
Candelina submexicana, 302
capsule expansion in Funaria, 457
Carex aquatilis, 215-
Carya laciniosa, 1-
Cassiope mertensiana, 314-; stelleriana, 437-; tetragona, 437-
Catagonium brevicaudatum, *157*, *162*; politum, 157, 162
Catascopium nigritum, 441, 482
Catharinaea. *See* Catharinea
Catharinea aequinoctialis, *161*, *162*; elamellosa, *160*, *162*; grossidens, *160*, *162*; integrifolia, *160*, *162*; nigricans, *160*, *162*; polycarpa, *160*, *162*; pseudo-polytrichum, *521*; pygmaea, *161*, *162*
Catillaria baeomma, *390*; biformigera, 390

CCC (2-chloroethyl-trimethylammonium chloride). effects on plastid ultrastructure in Marchantia, 22
Celtis reticulata, 269-; spp., 269-
Cenomyce stellaris, *363*, *391*
Cephalozia affinis, 430; bicuspidata, 185; bicuspidata subsp. ambigua, 185, 430; bicuspidata subsp. bicuspidata, 430; bicuspidata subsp. lammersiana, 185, 430; catenulata, 430; connivens, 430; lacinulata, 430; loitlesbergeri, 430; lunulifolia, 185, 430; macounii, 430; media, 1-; pleniceps, 185, 430; spp., 314-
Cephaloziaceae, 185, 314-, 430
Cephaloziella alpina, 430; arctica, 431; brinkmanii, 431; dentata, 431; divaricata, 186, 431; elachista, 431; hampeana, 431; phyllacantha, 431; rubella, 186, 431; sp., 55-, 266-; subdentata, 431; turneri, 431
Cephaloziellaceae, 186, 430
Ceratodon novogranatensis, 128; purpureus, 86-, 95-, 188-, 279, 339-, 441, 482; stenocarpus, 128, 269-
Cetraria ciliaris, 390; commixta, 448; coralligera, 302; cucullata, 448; delisei, 390, 449; ericetorum, 449; fendleri, 302; halei, *390*; hepatizon, 449; islandica, 449; laevigata, 449; merrillii, 390, 392; nivalis, 449; oakesiana, 390; pinastri, 449; richardsonii, 449; subalpina, 449; tilesii, 55-, 449
Chad [Thchad]. *See* Africa
Chamaenerion angustifolium, 172-
Chandonanthus setiformis, 183
Cheilolejeunea **intricata**, comb. nov., 514; **savatieriana**, comb. nov., 514
Cheilothela boliviana, 133, 162; chilensis, 133, 169; revoluta, 133, 162
CHILD, J. *See* ALLISON, K. W. & J. CHILD
Chile. new species and new combinations of hepatics from southern, 514
Chilopsis linearis, 269-
Chiloscyphus pallescens, 185, 429; pallescens var. fragilis, 429; polyanthus, 185, 429; polyanthus var. rivularis, 429
China. distribution of Rozea in, 406-
Chiodecton sanguineum, 358
chloroatranorin, 385-
chlorophyll and carotenoid production. in germinating Polytrichum spores, 332
chloroplasts. ultrastructure in Marchantia, 25*, 26*, 29*
Chorisodontium, 122-; germainii, 130, 163; mittenii, 130, 163; nigricans, 130, 163; speciosum, 130, 163; spectabile, 130, 163
Chrysoblastella boliviana, *133*, *162*; revoluta, *133*, *162*
CHURCHILL, S. P. Contributions Toward a Moss Flora of Nebraska, 241
Cinclidium arcticum, 502; stygium, 442; subrotundum, 442
Cirriphyllum andinum, 155; cirrosum, 442, 482, 502; laevifolium, 155
Cirsium spp., 172-
Cladia, 390
Cladina alpestris, *364*, 449; arbuscula, 449; impexa, 390; mitis, 391, 449; rangiferina, 391, 449; ***stellaris***, comb. nov., 363, 391; terrae-novae, 391; terrae-novae fo. **cinerescens**, comb. nov., 391
Cladonia, 266-; acuminata, 449; alpestris, *364*, *391*; amaurocraea, 449; bacillaris, 302; balfourii, 302; bellidiflora, 242-; bellidiflora fo. vestita, 242-; boryi, 391; botrytes, 449; cariosa, 449; carneola, 449; cenotea, 449; cervicornis, 172-; chlorophaea, 1-, 172-, 302, 450; clavulifera, 1-; coccifera, 172-, 450; coniocraea, 172-; cornuta, 450; cornuta β rei, *245*; cristatella, 1-; cyanipes, 450; ecmocyna, 450; fibula, *245*; fimbriata, 450; fimbriata *nemoxyna, *245*; floerkeana, 172-; furcata, 391, 450; gonecha, 450; gracilis, 450; gracilis fo. aspera, *245*; gracilis var. gracilis, 245; impexa, 172-; lepidota, 450; macilenta, *242-*, 391; macrophyllodes, 450; major, 450; mitis, *391*; multiformis fo. multiformis, 450; multiformis fo. subascypha, 391, 450; nemoxyna, *245*; norrlinii, 450; pacifica, *390*; phyllophora, 450; pityrea, 450; pocillum, 450; pyxidata, 172-, 302, 450; rangiferina, 1-, *391*; rangiformis, 172-; rei, 242; spp., 172-; squamosa, 450; stellaris, *364*; subcervicornis, 450; terrae-novae fo. cinerescens, *391*; uncialis, 172-, 451; verticillata, 451
Clapodiella, 314-
Clavaria argillacea, 172-
Clevea hyalina, 186, 433
Cleveaceae, 186, 433
Climacium americanum, 95-; dendroides, 95-, 442
Cnestrum alpestre, 321-, 503; schisti, 503
Coccocarpia cronia, 303
Coccotrema maritimum, 391; pocillarium, 392
Coenogonium interplexum, 358; interpositum, 358
collatolic acid, 352
Collema coccophorum, 303, 451; conglomeratum, 303; furfuraceum, 303; furfuraceum var. luzonense, 392; luzonense, *392*; luzonense fo. pruinosum, 392; glaucophthalmum var. glaucophthalmum, 359; tenax, 303; tuniforme, 303
Cololejeunea biddlecomiae, 1-
Colombia. Polytrichum ericoides in, 93
Colorado. additions to the bryophyte flora of, 215

connorstictic acid, 224-
Conocephalaceae, 186, 433
Conocephalum conicum, 1-, 186, 433
Conostomum, 122-; aequinoctiale, 144; aequinoctiale fo. breviseta, 144; australe, *144*, *162*; cleistocarpum, 144; macrotheca, 144; pentastichum, 144, 162; tetragonum, 442, 503
conpsoromic acid, 224-, 296-
constictic acid, 385-
Convolvulus arvensis, 172-
Cornicularia aculeata, 172-, 392, 451; californica, 390, 392; divergens, 451
Cosçinodon bolivianus, *137*, *162*; cribrosus, 482; trinervis, *137*, *162*
Costa Rica. Stenodictyon wrightii in, 84
Cratoneuron arcticum, 482, *483*; commutatum var. commutatum, 442; commutatum var. falcatum, 442; falcatum, 219; filicinum, 95-, 442, 482; filicinum var. **arcticum**, comb. & stat. nov., 483; filicinum var. curvicaule, 154, 164; filicinum var. julocladon, 154; oedogonium, *155*, *162*; punae, 154, 164; submersum, 154; williamsii, 442
Crosby, M. R. Recent Literature on Mosses, 107, 535
Crosby, M. R. The Correct Name for the Moss Duseniella, 246
Crosby, M. R. The Relationship of the Mosses Hookeria wrightii and Stenodictyon sericeum, 83
Crossidium aberrans, 282; chloronotos, 516; desertorum, 282; rotundifolium, 133; squamigerum, 517; wallisii, 133
Crum, H. *See* Buck, W. R. & H. Crum and Ireland, R. R. & H. Crum
Cryphaea boliviana, 148; brachycarpa, *148*, *162*; gracillima, 148; hygrophila, 148; hygrophila var. nitens, *148*, *162*; jamesonii, 148; latifolia, 148; macrospora, 148; microspora, 148, 163; nitens, 148, 162; patens, 148; patens var. dilatata, 148; pilifera, 148; ramosa, 148; ruficalyx, *148*, *163*; tenuicaulis, 148
Cryphaeaceae, 148
cryptochlorophaeic acid, 224-
Ctenidiadelphus cylindricarpus, 406-
Ctenidium malacodes, 159; plumulosum, 159; subimponens, *159*, *163*
Cukierski, M. *See* Paolillo, D. J. & M. Cukierski
Culberson, C. F. & T. L. Esslinger. 4-O-Methylolivetoric and Loxodellic Acids: New Depsides from New Species of Brown Parmeliae, 42
Culberson, C. F. *See also* Culberson, W. L. & C. F. Culberson
Culberson, W. L. Recent Literature on Lichens, 113, 523
Culberson, W. L. & C. F. Culberson. A Tribute to Yasuhiko Asahina, 258
cultivation of wind-blown bryophyte gametophyte fragments, 55
Cupressina entodonticarpa, *159*, *163*
Cupressus arizonica, 269-
Cyathodium acrotrichum, 234-; aureonitens, 234-; cavernarum, 234-; denticulatum, 234-; flabellatum, 234-; foetidissimum, 234-; smaragdinum, 234-; **tuberculatum**, sp. nov., 235, 236*; tuberosum, 234-
Cyclodictyon aeruginosum, 151; albicans, 151; angustirete, 151; bakeri, 151, 164; breve, 151; humile, 151; limbatum, 152; obscurum, 152; pandurifolium, 152; plicatulum, 152, 164; purpurascens, *152*, *163*; pusillum, 152; roridum, 152; stephanii, 152, 163; tocoraniense, 152; ulophyllum, 152
Cylindrothecium heteropterum, *415*; mandonii, *156*, *163*; subjulaceum, *416*
Cymbalaria muralis, 172-
Cynodontium gracilescens, 215
Cyphelium tigillare, 392
Cyrtomnium hymenophylloides, 442, 483; hymenophyllum, 442
cytokinin effect on bud formation in Microdus miquelianus, 495
cytoplasmic changes in spermatogenous cells of Polytrichum, 466

Dactylina arctica, 451; madreporiformis, 451; ramulosa, 451
Daltonia gracilis, 152, 163; hampeana, *152*, *163*; irrorata, *152*, *163*; jamesonii, 152; latolimbata, 152; lindigiana, 152, 163; longifolia, 152; minutifolia, *152*, *163*; pellucida, 152; pulvinata, 152; subirrorata, *152*, *163*; tenuifolia, 152; trachyodonta, 152, 163
Dasylirion leiophyllum, 269-
Delgadillo M., C. Indusiella andersonii sp. nov. in Northern Africa, 98
Delgadillo M., C. & W. D. Reese. Notes on Tropical Mosses of Mexico, 511
demethylnotatic acid, 224
Dendroligotrichum, 122-
Denmark. distribution of cryptogamic epiphytes in relation to air pollution and bark properties in, 86
depsides, 78-; in Parmeliae, 42
Dermatocarpon fluviatile, 451; hepaticum, 303, 451; lachneum, 296-; miniatum, 1-, 303; moulinsii, 303; tuckermanii, 303
Deschampsia flexuosa, 172-
Desmatodon cernuus, 85, 442, 483; convolutus, 282; guepinii, 282; heimii, 85, 442;

heimii var. arctica, 483; latifolius, 442, 503; leucostoma, 216, 442, 483, 503; plinthobius, 216, 241, 282; sp., 55-; systylius, 442, 503
Desmaziera cephalota, 364-; ceruchis, 364-
development. *See* growth
Dichelyma uncinatum, 218
Dichodontium pellucidum, 215-, 442, 503
Dicnemoloma kunertii, 130, 165
Dicranaceae, 128, 215, 279
Dicranella. *See also* Anisothecium and Microdus; angustifolia, 130; apolensis, 130; beyrichiana, *131*, *163*; beyrichii, *131*, *163*; boliviana, 130; callosa, 130; campylophylla, *128*, *163*; crispa, 442, 483; grevilliana, 442; guilleminiana, 130; heteromalla, 130, 241, 266-; hilariana, 131, 163; jamesonii, *128*, *163*; kunzeana, *131*, *163*; laxiretis, *128*, *163*; macrostoma, *128*, *163*; nanocarpa, 131, 161; perrottetii, 131; rufescens, 102; schreberiana, 216; submacrostoma, *128*, *163*; subserrulata, *131*, *163*; subulata, 442; tenuirostris, *131*, *163*; varia, 442
Dicranoloma, 122-
Dicranoweisia brunnea, 131; cirrata, 279; crispula, 442, 483, 503; fallax, *129*, *163*; flexipes, *129*, *163*
Dicranum, 122-; aciphyllum, 55-; acutifolium, 442, 483; angustatum, 442, 483; bolivianum, *131*, *163*; densicoma, *129*, *163*; elongatum, 442, 483; erectum, *130*, *163*; flaccidissimum, 131, 163; frigidum, 131; fulvum, 1-, 95-; germainii, *130*, *163*; groenlandicum, 503; laevidens, *483*; latifolium, 503; leucognodes, *129*, *163*; majus, 503; majus var. orthophyllum, 442; montanum, 95-; muehlenbeckii var. cirratum, 442; muehlenbeckii var. muehlenbeckii, 443; multicapsulare, *130*, *163*; nanofilifolium, *130*, *163*; nigricans, *130*, *163*; perexile, *130*, *163*; perreduncum, *130*, *163*; scoparium, 188, 199-, 201*, 443, 503; speciosum, *130*, *163*; spectabile, *130*, *163*; spiripes, *129*, *163*; spurio-concolor, *130*, *163*; tamarindifolium, *40*; triviale, *130*, *163*; undulatum, 188-, 443; viridulum, *40*
Didymodon aeneus, *229*; amblystegius, *133*, *163*; angustifolius, *134*, *163*; angustulus, 134, 163; argentinicus, 134, 163, 169; asperifolius, 55-, 443, 483, 503; campylopyxis, *133*, *163*; contortus, 134; decolorans, *133*, *163*; decolorans var. brevifolia, *133*; decolorans var. obtusus, *133*; jamesonii, *133*, *163*; johansenii, 443, 483, 503; luteus, *134*, *163*; macrophyllus, 134; merceyoides, 134; mexicanus var. subulatus, 283; pelichucensis, 134; perexilis, 134, 161; recurvirostris, *229*; reedii, 216; rigidulus, 215-; rubiginosus, *134*, *163*; semivaginatus, 134, 169; subtophaceus, 134; tophaceus, 134, 169, 216, 283
diffractaic acid, 385-
Digitalis purpurea, 172-
Dilaenaceae, 186, 432
Dimerella pyrophthalma, 359
Diospyros texana, 269-
Diphyscium foliosum, 1-
Diplophyllum albicans, 428; apiculatum, 1-; obtusifolium, 428; plicatum, 428; taxifolium, 184, 428
Diploschistes scruposus, 451
Dirinaria applanata, 359; aspera, 304; frostii, 304
Dissodon andinus, 152; bolivianus, 152
Distichium capillaceum, 128, 443; capillaceum var. strictifolium, 128, 163; hagenii, 483, 503; inclinatum, 85, 443, 483; sp. 55-, 61*; strictifolium, *128*, *163*
distribution. *See* phytogeography
Ditrichaceae, 128, 278
Ditrichum bogotense, 128; capillare, 128, 165; crinale, 128; flexicaule, 55-, 443, 483; levieri, 128, 167; pallidum, 1-, 101-; rhynchostegium, 101; rufescens, 128; subflexuosum, 128, 165; submersum, 128
DOUGLAS, G. W. & D. H. VITT. Moss-Lichen Flora of St. Elias-Kluane Ranges, Southwestern Yukon, 437
Drepanocladus aduncus, 443, 483; aduncus subsp. barbeyi, 154, 164; badius, 443; brevifolius, 208-; capillifolius, 219; exannulatus, 47-, 154, 443; exannulatus fo. **pseudotrichophyllus**, nom. nov., 219; exannulatus var. rotae, *219*; fluitans, 219; revolvens, 208-, 443, 483; sendtneri, 154, 443; sp., 55-, 154; trichophyllus, 215-; uncinatus, 47-, 55-, 443; vernicosus, 443
Dryas drummondii, 437-; octopetala, 437-
Dryopteris marginalis, 222-
Duseniella, *247*; genuflexa, *247*

ecology. bryophytes used in construction of bird nests, 95; caloric values of three Alaskan-Arctic mosses, 76; distribution of cryptogamic epiphytes in relation to air pollution and bark factors, 86; effect of gamma radiation on regeneration in leaf and stem-leaf cultures of Polytrichum, 488; effect of light and other factors on capsule expansion in Funaria, 457; effects of pulp mill emissions on lichens, 248; effects of substrate and microtopography on vegetative propogation of bryophytes, 199; growth in culture of wind-blown bryophyte gametophyte fragments, 55; lichen ecology of metal spoil tips: effects of metal ions on ascospore viability, 172;

Ramalina leptocarpha and R. subleptocarpha, a fertile-sorediate species pair, 364; relationship of terrestrial mosses to forested flood plains, 64; saxicolous bryophyte and macrolichen associations, 1; seasonal growth of Parmelia caperata, 360; sensitivity of lichens to nitrogen dioxide fumigations, 103; some mosses from Kuwait, 515
Ecklonia radiata, 22-
Ectropothecium aeruginosum, 159; amphibolum, *160*, *163*; apiculatum, *159*, *163*; campanulatum, 159; effusum, 159; globitheca, 159, 163; vesicularis, *160*, *163*
EGAN, R. S. New and Additional Lichen Records from Texas, 224
EGAN, R. S. Speerschneidera euploca, 579*
EGAN, R. S. (review) S. R. Rushforth. The Plant Kingdom, 549
electron micrographs. germinating Dicranum spores, 190*, 191*, 193*, 194*, 195*, 197*; antheridia of Polytrichum, 468*, 470*, 472*, 474*, 476*, 478*; cells and chloroplasts of Marchantia, 24*, 25*, 26*, 29*; leaf sections of Ceratodon purpureus, 343*; leaf sections of Hygrohypnum luridum, 345*; leaf sections of Hypnum curvifolium, 344*; leaf sections of Mnium cuspidatum, 343*; leaf sections of Mnium punctatum, 345*; leaf sections of Tortula ruralis, 343*; protonemata of Ceratodon purpureus, 347*; protonemata of Polytrichum commune, 336*, 337*
Ellesmere Island. *See* Canada
EM. *See* electron micrographs
Empretrum hermaphroditum, 47-; nigrum, 437-
Encalypta. Key to gymnostomous spp. in North America, 322; affinis, 443, 503; alpina, 321-, 443, 483, 503; asperifolia, 132, 163; brevicolla, 321-, 503, brevipes, 322, 323*, 327*, 501; distribution in North America, 328*; ciliata, 321-, 503; coarctata, 132; leiotheca, 132; longicolla, 321-, 503; mutica, 325*, 326, 327*, 501, distribution in North America, 328*; procera, 216, 321-, 443, 483; rhaptocarpa, 321-, 443, 483; vernicosa, *132*, *163*; vulgaris, 279, 321-, 444; vulgaris fo. pilifera, 321-
Encalyptaceae, 132, 216, 279
ENGEL, J. J. Austral Hepaticae VI. Some New Species and New Combinations of Taxa from Southern Chile, 514
Entodon, 406-; beyrichii, 156, 163; bolivianus, 156, 164; cladorrhizans, 95-, 102; concinnus, 444; erythropus, *156*, *163*, 293; flavissimus, 156; flexipes, 156; germainii, *156*, *163*; gracilisetus, 156; hampeanus, 156; jamesonii, 156, 163; lutescens, 156; macropodus, 156, 163; mandonii, 156, 163; microcarpus, 156; nanoclimacium, 156; nanoclimacium var. macropterus, 156; pallidisetus, 156; polycarpus, *156*, *163*; polysetus, *156*, *163*; seductrix, 95-; suberythropus, 156; viridis, *156*, *163*
Entodontaceae, 156, 220, 293, 406-
Entosthodon acidotus, *138*, *163*; altisetus, *138*, *163*; apiahyensis, *138*, *163*; apiculatus, 138; cartilagineus, *138*, *163*; faucium, *138*, *163*; fontanus, *138*, *163*; glabripes, *138*, *163*; lindigii, *138*, *163*; mandonii, 138; papillosus, *138*, *163*; sipascoyae, *138*, *163*; sonorae, 287; subtilis, *138*, *163*; verrucosus, *138*, *163*
Ephebe lanata, 392
Epipterygium mandonii, 140, 162; pellucens, 140
Equisetum, 188-
Eriophorum angustifolium, 47-
Eriopus papillatus, 152
Erpodiaceae, 144
Erpodium balansae, *144*, *163*; beccarii, 144; coronatum, 144; glaziovii, 144, 163
Erythrina crita-galli, 358-
Erythrodontium brasiliense, 156, 163; brasiliense fo. **brevipes**, comb. nov., 156, 163; brasiliense var. brevifolium, 156; germainii, *156*, *163*; germainii var. brevipes, *156*, *163*; longisetum, 156; macrocarpum, 156; squarrosum, 156
Erythrophyllopsis boliviana, *134*, *163*; challaensis, 134, 169; fuscula, 134, 163
Erythrophyllum decolorans, *133*, *163*; rubiginosum, *134*, *163*
ESSLINGER, T. L. *See* CULBERSON, C. F. & T. L. ESSLINGER
Euphorbia antisyphilitica, 269-
Eurhynchiella semitorta, 155, 167; toncolensis, 155, 167
Eurhynchium exasperatum, *156*, *163*; hians, 64-, 95-; integrifolium, 155; oedogonium, 155, 162; pulchellum, 95-, 293, 444, 484; subdenticulatum, 155
Eustichia spruceana, 142
Eustichiaceae, 142
Evernia divaricata, 451; esorediosa, 451; mesomorpha, 392; perfragilis, 451
exsiccata. Lichenes Canadenses, 385

Fabronia andina, 153; attaleae, 153; balansae, 153; ciliaris, 102, 241, 291; podocarpi, 153; polycarpa, 153; pusilla, 291; seligeriaceae, 153; singulidens, 153; wrightii, 291
Fabroniaceae, 153, 291
Fagus grandiflora, 1-; sylvatica, 172-
Felipponea montevidensis, 148, 165

Festuca altaica, 437-; ovina, 172-; rubra, 172-
FIFE, A. J. *See* BRASSARD, G. R., D. P. WEBER & A. J. FIFE
Fissidens, 227-. *See also* Moenkemeyera & Simplicidens; sect. Bryoidium, 35-; sect. Semilimbatum, 35-; sect. Semilimbatus, 35-; subsect. Semilimbatus, 35-; adianthoides, 503; amboroicus, 127; andersonii, *40*; arcticus, 321-, 503; asplenioides, 127; bambergeri, 35-; bockii, 127; bryoides, 35, 102, 215, 278, 444, 511; bryoides var. brevifolius, *41*; bryoides var. incurvus, *40*; burelae, 127; bushii, 278; carionis, *128*, *163*; crispus, 127; elegans, 511; excurrentinervis, 127; exiguus, *40*; exiguus var. falcatulus, 38*, *40*; falcatulus, *40*; fontanus, 278; gymnandrus, 35-; hornschuchii, *128*, *163*; incacorralis, 128; incisus, 128; inconstans, 35-; incurvus, *40*; incurvus var. brevifolius, *40*; innovans, 128; kegelianus, 128; limbatus, 278; macroblastus, 128; macrophyllus, 128; mateoënsis, 128; minutulus, 1-, *40*; mollis, 512; obtusifolius, 241, 516; obtusifolius var. apiculatus, 278; obtusifolius var. obtusifolius, 278; oligophyllus, 128, 163; osmundioides, 1-, 503; pauper, 128; prionodes, 128, 163; repandus, 128, 163, 278; revolvens, 128; rigidulus, 128; sanctaecrucis, 128; stenosemus, *128*, *163*; subbasilaris, 1-; sublimbatus, 215, 278; synoicus, 35-; taxifolius, 1-, 64-; terebrifolius, 128; texanus, *40*; tocoraniensis, 128; viridulus, *40*, 85, 106-; viridulus var. brevifolius, 38*, *41*; viridulus var. tamarindifolius, *40*; viridulus var. texanus, *40*; wallisii, 128
Fissidentaceae, 127, 215, 277
Flabellidium spinosum, 155
Floribundaria flaccida, *149*, *163*; tenuissima, *149*, *164*
floristics. *See* phytogeography
Fontinalaceae, 218
Fontinalis novae-angliae, 95-; turfacea, *155*, *164*
Forsstroemia coronata, 148; producta, 148, 162
Fossombronia, 466-
Fragaria vesca, 172-
Fraxinus excelsior, 86-, 172-
free amino acids in lichens, 354
FRENCH, J. C. & D. J. PAOLILLO. Effect of Light and Other Factors on Capsule Expansion in Funaria hygrometrica, 457
Frullania asagrayana, 1-, 95-; eboracensis, 95-; riparia, 1-
Fucus vesiculosus, 22-
FULFORD, M. Recent Literature on Hepatics, 253, 370, 531
Fulgensia bracteata, 451
fumarprotocetraric acid, 242-, 296-, 385-
Funaria, 466-; acidota, 138, 163; acutifolia, 138; altiseta, 138, 163; americana, 287; andicola, 138; apiculatipilosa, 287; arctica, 484, 523; boliviana, 138, 164; calvescens, *138*, *164*; cartilaginea, 138, 163; faucium, 138, 163; fontana, 138, 163; glabripes, 138, 163; hygrometrica, 124*, 138, 287, 444, 457, 463*, 464*, 515; hygrometrica var. calvescens, 138, 164; hygrometricoides, *138*, *164*; incurvifolia, 138; inflata, 138; lindigii, 138, 163; linearidens, 138; macrospora, 138; meeseacea, 138; papillosa, 138, 163, 164; polaris, 503; riparia, 138, 163; sipascoyae, 138, 163; soratensis, *138*, *164*; sp., 95-; subtilis, 138, 163; tucumanica, *138*, *164*; verrucosa, 138, 163, 164
Funariaceae, 138, 287
Fuscidea mollis, 392

Galax aphylla, 222-
GALLOWAY, D. J. William Martin (1886–1975), 374
GELLERMAN, J. L. *See* SWANSON, E. S. et al.
Geocalyx graveolens, 185, 268*, 429
Geothallus, 314-
Gertrudia validinervis, *134*, *164*; validinervis var. serrato-pungens, *134*, *164*
Gertrudiella ferruginea, 134, 169; validinervis, 134, 164
gibberellin, effects on plastid ultrastructure in Marchantia, 22
Globulina boliviana, *133*, *164*
Globulinella globifera, 283; peruviana, 229
glomellic acid, 42
glomelliferic acid, 42
Glyphomitrium cochabambae, *144*, *164*; ferrugineum, *145*, *164*; papillosum, *145*, *164*; sellowianum, *145*, *164*
GODFREY, J. D. Schofieldia, a New Hepatic from the Pacific Northwest, 314
Gonohymena sp., 304
GRADSTEIN, S. R. A Taxonomic Monograph of the Genus Acrolejeunea (review), 380
Graphis lineola var. marginata, 359; pavoniana, 359; scripta, 176, 359; spp., 172-
GREENE, S. W., B. G. BELL, G. C. S. CLARKE & M. E. NEWTON. A synoptic Flora of South Georgian Mosses (review), 122
Greenland. distribution of Grimmia cribrosa in, 356, 356*; Schistidium holmenianum in, 208; Sphagnum pylaesii in, 47
GRIFFIN, D. *See* W. D. REESE & D. GRIFFIN
Grimaldiaceae, 186, 432
Grimmia, 208-, 406-. *See also* Coscinodon & Schistidium; affinis, 286, 444, 503; alpicola, 286, 444, *485*; andina, 136, 168;

andreaeacea, 136; angustifolia, 137, 168; anodon, 444, 482; anomala, 217; apocarpa, 1-, 137, 168, 286, *485*; apocarpa var. apocarpa, 444, 504; apocarpa var. nigrescens, 208-, 485; apocarpa var. stricta, 444, *485*, 504; arizonae, 286; atricha, 217; bicolor, 137; **boliviana**, comb. nov., 137, 162; brachypus, *137*, *164*; calycina, 137; chocayae, 137, 168; cribosa, 286, 355, *482*; distribution in North America of, 356*, crispula, *137*, *164*; dimorpha, *137*, *164*; dupretii, 217; elongata, 217; flaccida, 504; flexicaulis, 137; fontana, 137, 168; herzogii, 137; leucophaeola, 137; longirostris, 137; malacophylla, 137, 168; micromitria, 137; micro-ovata, 137; mollis, 208-; nanoglobosa, 137; navicularis, 137; nigella, 137; olivaceae, 137; olneyi, 1-; ovalis, 286; pansa, 137; praemorsa, 137, 168; quadricruris, 137; quatricruris, *137*; sp., 55-; speirophylla, 137; squamatula, 137; streptophylla, 137, 168; subovata, 137; subpraemorsa, 137, 168; subquadricruris, 137; subquatricruris, *137*; tenera, *485*; torquata, 321-; trichophylla, 85; trichophylloidea, 137; trinervis, 137, 162; tristicha, 137; tristicha var. comosa, 137; tunariensis, 137, 168; wrightii, 286, 355-
Grimmiaceae, 98-, 136, 217, 286
GROSSMAN, H. H. & C. J. HILLSON. The Effect of Chronic Low-Level Gamma Radiation on Polytrichum commune. II. Regeneration in Leaf and Stem-Leaf Cuttings, 488
Groutiella mucronifolia, 145, 165
growth. cultures of wind-blown bryophyte gametophyte fragments, 55; cytokinin induced bud formation in Microdus miquelianus, 495; effect of gamma radiation on regeneration in leaf and stem-leaf cuttings of Polytrichum, 488; effect of metal ions on ascospore viability, 172; effects of gibberellin and growth retardants on plastid ultrastructure in Marchantia, 22; generation of Fissidens taxifolius, 72*; germinating Polytrichum spores, 332; in spermatogenous cells of Polytrichum, 466; of capsules in Funaria, 457, 463*, 464*; of Parmelia caperata, 360
Guadeloupe. *See* West Indies
Guatemala. notes on Pottiaceae in Middle America, 227; distribution of Rozea in, 406-
Guignardia alaskana, 396, 397*; prasiolae, 397; -Prasiola complex, 400*, 401*; ulvae, 397
Gymnocolea inflata, 183
Gymnodinium breve, 22-
Gymnomitriaceae, 184, 428
Gymnomitrion concinnatum, 428; corallioides, 184, 428; obtusum, 428; pacificum, 428; sphacelata, 428
Gymnostomum aeruginosum, 1-, 85, 216, 283; calcareum, 516; phascoides, *384*; recurvirostre, 215-, 283, 444, 484; valerianum, 229
gyrophoric acid, 43, 78-, 224-
Gyrothyra, 314-
Gyroweisia boliviana, *133*, *164*

Haematomma lapponicum, 451
HALE, M. E. Hypotrachyna Showmanii, a New Lichen from Eastern North America, 78
Haplocarpon crustulatum, 393; macrocarpum, 393
Haplocladium amblyostomum, *154*, *164*; austro-serpens, *154*, *164*; laterculi, *154*, *164*; microphyllum, 154, 164, 292
Haplodontium crassinervium, 140; crassinervum, *140*; cuspidatum, 140; herzogii, 140; humipetens, 140; jamesonii, *140*, *164*; megalocarpum, 140, 164; microphyllum, 102; sanguinolentum, 140; subsplendidum, 140; vilocense, 140
Harpanthus flotowianus, 185
Harpolejeunea intricata, *514*
Harrisonia humboldtii, *148*, *164*; mandonii, *147*, *148*, *164*
Hawaii. Ramalina farinacea, in, 499
HAYNES, J. D. Botany. An Introductory Survey of the Plant Kingdom (review), 264
Hechtea scariosa, 269-
Hedera helix, 172-
Hedwigia albicans, *147*, *164*; ciliata, 1-, 95-, 147, 164, 291
Hedwigiaceae, 147, 290
Hedwigidium imberbe, *147*, *164*; imberbe var. macrocalyx, *147*, *164*; integrifolium, 147, 162, 164
HEGEWALD, E. *See* ZANDER, R. H. & E. HEGEWALD
Helicoblepharum venustum, 152
Helicodontium capillare, 153; rhyparobolax, 153; sp., 153, 167; spicatinervium, *155*, *164*; tenuirostre, 153
Helicophyllaceae, 147
Helicophyllum torquatum, 147
Helodium paludosum, 101
hepatic ecology. *See* ecology
Heppia lutosa, 304
Heracleum sphondylium, 172-
HERMANN, F. J. Additions to the Bryophyte Flora of Colorado, 215
HERMANN, F. J. Recopilación de los Musgos de Bolivia, 125
Herpetineuron toccoae, 292

Herzogiella boliviana, 157, 166; turfacea, 95-
Heterocladium dimorphum, 444
Heterodermia appalachensis, 304; diademata, 304; domingensis, 304; granulifera, 304; leucomela, 304; rugulosa, 304
Heuchera villosa, 222-
Hicks, M. L. Lophozia heterocolpa, New Disjunct for North Carolina, 222
Hieracium spp., 172-
Hilaria mutica, 269-
Hillson, C. J. *See* Grossman, H. H. & C. J. Hillson
HIMT (4-hydroxy-5-isopropyl-2-methylphenyl trimethylammonium chloride-1-piperidine carboxylate). effects on plastid ultrastructure in Marchantia, 22
Holcus spp., 172-
Holomitrium aristatum, 131; bolivianum, *135*, *164*; crispulum, 131; macrocarpum, *135*, *164*
Homalia glabella, 511
Homalotheciella subcapillata, 95-
Homalothecium, 406-; sericeum, 86-
Homomallium adnatum, 1-; mexicanum, 220, 294
homosekikaic acid, 242-
Hong, W. S. Annotated Checklist of the Hepatics of Idaho, 422
Hong, W. S. & D. H. Vitt. A List of Hepaticae from West-Central Alberta, 180
Hookeria sect. Hypnella, 83-; subsect. Hylotapis, 83-; acutifolia, 152; ancistrodes, *247*; bakeri, *151*, *164*; brunneophylla, *152*, *164*; (Chaetephora) incurva, 246-; crispa, *152*, *164*; curviramea, *153*, *164*; falcata, *152*, *164*; integrifolia, *151*, *164*; pallidonitens, *153*, *164*; papillidioides, *152*, *164*; plicatula, *152*, *164*; (Pterygoph.) splendidissima, 246-; (Pterigophyllum) obscura, 246-; purpureophylla, *152*, *164*; scabripes, *151*, *164*; sigmatelloides, *152*, *164*; undatula, *152*, *164*; wrightii, *83*
Hookeriaceae, 151
Hookeriopsis asprella, 152; brunneophylla, 152, 164; crispa, 152, 164; falcata, 152, 164; falcata var. latifolia, 152; incurva, 152; langsdorffii, *164*; lepidopiloides, 152; longiseta, 152; pachydictyon, 152; papillidioides, 152, 164; purpureophylla, 152, 164; subsecunda, 152; undatula, 152, 164; variabilis, 152; williamsii, 152
Horton, D. G. & B. M. Murray. Encalypta brevipes and E. mutica, Gymnostomous Species New to North America, 321
Husnotiella glossophylla, 134; revoluta, 216, 283; torquescens, 283
Hydropogonaceae, 147
Hydropogonella gymnostoma, 147
Hygroamblystegium curvicaule, *154*, *164*; fluviatile, 219; orthocladon, 219; punae, *154*, *164*; punae var. tenuinerve, *154*, *164*; tenax, 95-
Hygrobiella, 314-; laxifolia, 430
Hygrodicranum bolivianum, 131
Hygrohypnum aureum, 154; bestii, 219, 504; cochlearifolium, 504; eugyrium, 95-; luridum, 339-, 444, 484; ochraceum, 484; pelichucense, 155, 167; polare, 484, 504; smithii, 219; validum, 155
Hylocomium pyrenaicum, 504; splendens, 444; splendens var. obtusifolium, 484
Hylotapis, 83-
Hymenostomum anomalum, 134; rostellatum var. phascoides, *384*; tortile, 283
Hymenostylium contextum, 134; recurvirostre, 16-
Hyophila contermina, *134*, *164*; involuta, 134, 164, 283; involutifolia, 134; minutissima, 134; tortula, *134*, *164*, 512
Hypnaceae, 159, 220, 293, 406-
Hypnella brotheri, 152; pilifera, 152; sigmatelloides, 152, 164; wrightii, *84*
Hypnum amabile, 159, 162, 163; andrieuxii, *413*; bambergeri, 208-, 321-, 444; barbeyi, *154*, *164*; bolivianum, *156*, *164*; callichroum, 504; cirrhato-revolvens, 159; "cristatum," 95-; cupressiforme, 86-, 159, 199-, 208-, 222-, 294, 444, 484; cupressiforme var. uncinulatum, 159, 164; curvifolium, 95-, 339-; entodonticarpum, *159*, *164*; fertile, 95-; fulvum, *419*; imponens, 95-; latifolium, *159*, *164*; lindbergii, 95-, 444; mandonii, 159; pallescens, 95-; plicatulum, 444; procerrimum, 444, 484; protensum, 159, 165; pterogonioides, *417*; recurvatum, 504; revolutum, 444, 484; spiripes, 159, 163, 164, 168; subimponens, 484; turgescens, 159; vaucheri, 294, 444
Hypogymnia atrofusca, 451; austerodes, 451; enteromorpha, 248; physodes, 451
hypoprotocetraric acid, 224
Hypopterygiaceae, 153
Hypopterygium argentinicum, 153; bolivianum, 153; tamarisci, *153*, *164*; tamariscinum, 153, 164
Hypotrachyna formosana, 78-, 224-; livida, 224-; pustulifera, 78-, 225; **showmanii**, sp. nov., 78, 79*

Icmadophila ericetorum, 451
Idaho. annotated checklish of hepatics of, 422; Sphagnum in, 508
Illinois. Buxbaumia aphylla in, 266; saxicolous bryophyte and macrolichen associations in southern, 1
Impatiens pallida, 64-
India. a new Cyathodium from, 234; distribution of Rozea in, 406-

Indusiella **andersonii**, sp. nov., 99, 99*; thianschanica, 98-
Ionaspis melanocarpa, 393
IRELAND, R. R. Alar Cells of Rhacomitrium lawtonae, 238
IRELAND, R. R. & H. CRUM. Distribution of Grimmia cribrosa in North America, 355
irradiation. effect of gamma radiation on regeneration in leaf and stem-leaf cuttings of Polytrichum, 488
Isopterygium brachyneuron, *157*, *164*; chrismarii, 157; cylindraceum, 157, 168; distichaceum, 95-; elegans, 1-; homomallifolium, 80-; leucophyllum, 157; peruvianum, 157; pulchellum, 321-, 444; stigmocarpum, 157, 168; subglobosum, 157; tenerum, 157, 164; vagans, 157
Isothecium andrieuxii, *413*

Jaegerina scariosa, 149, 164
Jaegerinopsis brasiliensis, *149*, *164*
Jamesoniella autumnalis, 1-, 184, 427
JOHNSON, I. & U. SØCHTING. Distribution of Cryptogamic Epiphytes in a Danish City in Relation to Air Pollution and Bark Properties, 86
JUÁREZ, G., L. G. & A. J. SHARP. Weissia rostellata var. phascoides New to Mexico, 384
Jubula pennsylvania, 1-
Juglans nigra, 269-
Juncus squarrosus, 47-
Jungermannia (Lejeunea) intricata, *514*; lanceolata, 427; oligophylla, *514*; tubulata, *514*
Jungermanniaceae, 184, 427
Jungermanniales, 424
Juniperus deppeana, 269-; flaccida, 269-; pinchottii, 269-; spp., 269-; virginiana, 1-
Juratzkaea, 406-
JÜRGING, P. Epiphytische Flechten als Bioindikatoren der Luftverunreinigung-Dargestellt an Untersuchungen und Beobachtungen in Bayern (review), 264

Kalmia latifolia, 222-
Kansas. notes on mosses of, 101
KARUNEN, P., N. VALANNE & R. E. WILKINSON. Influence of S-Ethyl dipropylthiocarbamate on Growth, Chlorophyll and Carotenoid Production and Chloroplast Ultrastructure of Germinating Polytrichum commune Spores, 332
KAUSTINEN, H. *See* ALBERTINE, K. H. et al.
KERR, R. *See* SHERIDAN, R. P. et al.
Kiaeria glacialis, 504; starkei, 216, 504
KIRKPATRICK, R. *See* MAGILL, R. E. & R. KIRKPATRICK
KROG, H. & T. D. V. SWINSCOW. Pyxine azorea Nyl. is Parmelia caroliniana Nyl., 506
KUC, M. Bryogeography of Expedition Area, Axel Heiberg Island, N.W.T., Canada (review), 547
Kurzia **mollis**, comb. nov., 514; **setiformis**, comb. nov., 514; sylvatica, 1-
Kuwait. some mosses from, 515

Laccaria laccata, 172-
Laestadia prasiolae, 396
LANGE, B. Sphagnum pylaesii in Greenland, 47
Laportea canadensis, 64-
Larrea divaricata, 269-
Lasallia papulosa, 304
lateral archegonia in mosses, 229
LAYSER, E. F. *See* ANDRUS, R. E. & E. F. LAYSER
LEBLANC, F. (review) P. JÜRGING, Epiphytische Flechten als Bioindikatoren der Luftverunreinigung-Dargestellt an Untersuchungen und Beobachtungen in Bayern, 264
Lecanactis amylacea, 393; illecebrosa, *393*; illecebrosa var. megaspora, *393*; megaspora, 393
Lecanora sect. Petrasterion, 350-; atra, 350-, 451; beringii, 395; caesiocinerea, 451; calcarea, 386, 388*; candida, 451; cenisia, 451; chlarotera, 86-; christoi, 350-; chrysoleuca, 103-, 452; cinereofusca, 393, distribution of, 395*; **collatolica**, sp. nov., 351*, 352; conizaeoides, 86-; degelii, 393; dispersa, 86-; epibryon, 452; expallens, 394; frustulosa, 452; hageni, 452; laxula, 394; marina, *389*; melanophthalma, 452; muralis, 452; mutabilis, 399; pacifica, 394; palanderi, 395; polytropa, 452; rupicola, 395, 452; subfusca, 394; subfusca var. tumescens, 393; subrubra var. microcheila, 394; subrubra var. subrubra, 394; urceolaria, 399; verrucosa, 452; wisconsinensis, 452
lecanoric acid, 78-, 224-
Lecidea albofuscescens, 395; armeniaca, 452; atrobrunnea, 452; atromarginata, 452; berengeriana, 452; biformigera, *390*; crustulata, 393; cuprea, 452; decipiens, 452; elata, 396; flavocaerulescens, 451; fuscella, 359; glomerulosa, 452; granulosa, 396, 452; lapicida, 452; limitata, 172-; macrocarpa, 172-, *393*; marginata, 396, 452; mollis, *392*; pantherina, 452; ramulosa, 396; rivulosa *β.* mollis, *392*; rubiformis, 452; subsorediza, 452; tessellata, 453
Lecideaceae, 350-
Lecidella eleochroma, 395; stigmatea, 453
Ledum groenlandicum, 47-
Leersia mutica, *326*
Leiomela bartramioides, 144, 161; brachy-

phylla, 144, 161; deciduifolia, 144; setifolia, 144, 161
Lejeunea lamacerina ssp. gemminata, 1-; savatieriana, *514*
Lembophyllaceae, 151
Lembophyllum bolivianum, *157*, *164*
Lepidopilidium synoicum, 152
Lepidopilum angustifrons, 152; auriculatum, 152; ballivianii, 152; brachyphyllum, 152; crispum, 153; curvirameum, 153, 164; filosum, 153; gertrudis, 153; herzogii, 153; intermedium, 153; malachiticum, 153; muelleri, 153; nanothecium, 153; ovatifolium, 153; pallido-nitens, 153, 164; tenuissimum, 153
Lepidozia filamentosa, 425; mollis, *514*; reptans, 182, 425; sandvicensis, 425; seriatitexta, *514*; setiformis, *514*
Lepidoziaceae, 182, 425
Lepraria membranacea, 1-; neglecta, 453
Leptobryum pyriforme, 199-, 444, 515; stellatum, 140, 169
Leptodictyum brevipes, 95-; loeskei, 155, 161; riparium, 95-; trichopodium, 64-, 95-, 219
Leptodontiella, gen. nov., 16; and Leptodontium, key to, 19; **apiculata**, comb. nov., 16
Leptodontium sect. Coronopapillata, 16-; acutifolium, *135*, *164*; acutifolium var. grimmioides, *135*, *164*, 169; albo-vaginatum, *133*, *164*; apiculatum, *16*; araucarieti, 16-, 134; brachyphyllum, 20, 134; calymperoides, *20*; capituligerum, 21, 134, 163, 164, 168, 169; cirrhifolium, *135*, *164*; erythroneuron, 134; filescens, *134*, *164*; filicola, 134; flexifolium, 134, 164, 165, 283; geheebiaeopsis, *134*, *164*; gracile, *134*, *164*; gracilescens, *134*, *164*; grimmioides, *135*, *165*; longicaule, 134, 165, 169; longicaule var. longicaule, 21; longicaule var. microruncinatum, 21, 134, 164, 165; luteum, 21, 134; mandonii, *134*, *165*; papillosum, *134*, *165*; planifolium, 135; proliferum, 135; pungens, 21, 135, 164, 165; rhacomitrioides, *135*, *165*; rufescens, *134*, *165*; spongiosum, *134*, *165*; sulphureum, *135*, *165*; **tricolor**, comb. nov., 20; turgidum, *135*, *165*; undulatum, *134*, *165*; undulatum fo. propagulifera, *134*, *165*; vaginatum, *135*, *165*; valerianum, 229; viticulosoides, 16-, 135, 164, 165; viticulosoides var. panamense, 135, 164, 165; viticulosoides var. viticulosoides, 21; wallisii, 21, 135, 165
Leptogiopsis complicatula, 397
Leptogium arsenei, 305; burnetiae var. hirsutum, 396; cyanescens, 359; denticulatum, 305; furfuraceum, 305; hirsutum, *396*; lichenoides, 1-, 305; milligranum, 305; phyllocarpum, 359; saturninum, 396
Leptohymenium chryseum, *411*; subjulaceum, *415*
Leptopterigynandrum austro-alpinum, 154
Leptotheca, 122-; boliviana, 142
Leptotrichum bolivianum, *128*, *165*; capillare, *128*, *165*; subflexuosum, *128*, *165*
Lepyrodon tomentosus, 148; tomentosus var. tunariensis, 148, 165; tunariensis, *148*, *165*
Lepyrodontaceae, 148
Leskea aciculata, *149*, *165*; boliviana, *153*, *165*; catenularia, 153, 167; fulva, 409, *419*; gracilescens, 1-; julicaulis, 153; plumaria, *154*, *165*; polycarpa, 86-; pterogonioides, *417*
Leskeaceae, 153, 218, 291
Leskeadelphus catenulatulus, 154, 167
Leskeodon andicola, 153
Letharia vulpina, 248, 396
Leucobryaceae, 131
Leucobryum albidum, 1-, 131, 165; calycinum, 131; crispum, 131; extensum, 131; fragile, 131; giganteum, 131; glaucum, 1-; linealifolium, *132*; linearifolium, 132; longifolium, *131*, *165*; macro-falcatum, 132; martianum, 132; minus, *131*, *165*; strictum, 132; subandinum, 132; tumidum, 132
Leucodon brachypus, 95-, 291; squarrosus, *148*, *165*
Leucodontaceae, 148, 291
Leucolejeunea unciloba, 1-
Leucoloma kunertii, *130*, *165*; macrodon, 131
Leucomiaceae, 153
Leucomium strumosum, 153
lichen ecology. *See* ecology
Lichen atrata, 389; fibula, 242-; rangiferinus alpestris, *364*
lichexanthone, 78-, 224-
light. effect on capsule expansion in Funaria, 457
Ligustrum lucidum, 358-
Limnobium protensum, *159*, *165*
Lindbergia brachyptera, 102, 241, 292; mexicana, 154, 292
Lindera benzoin, 1-
Lindigia acicularis, *149*, *165*; aciculata, 149, 165; debilis, 149
lipid composition of mosses, 339
Liquidambar, 511-; styraciflua, 1-
Liriodendron tulipifera, 1-
literature on hepatics, 253, 370, 531
literature on lichens, 113, 523
literature on mosses, 107, 535
Little Grand Canyon, Illinois, 1
lividic acid complex, 78-
Lobaria erosa, 224-; linita, 453; ravenelii, 224-

Lophocolea bidentata, 199-, 429; cuspidata, 429; heterophylla, 1-, 95-, 185, 430; minor, 1-, 185, 266-, 430
Lophocoleaceae, 185, 429
Lophozia alpestris, 183, 426; ascendens, 183, 220, 426; atlantica, 427; attenuata, 427; badensis, 426; bantriensis, 183, 426; barbata, 183, 425; binsteadii, 427; collaris, 183, 426; excisa, 183, 426; floerkei, 184, 199-, 427; gillmani, 183, 426; grandiretis, 184; hatcheri, 183, 426; heterocolpa, 183, 222, 426; incisa, 184, 426; kunzeana, 184, 427; longidens, 183, 426; lycopodioides, 183, 426; obtusa, 427; opacifolia, 314-; porphyroleuca, 183, 426; quadriloba, 184, 427; rutheana, 183, 426; sp., 55-; ventricosa, 183, 426; ventricosa var. longiflora, 184, 426; ventricosa var. silvicola, 184, 426; wenzelii, 314-, 426
Lophoziaceae, 183, 220, 314-, 425
loxodellic acid, 42
Lunularia cruciata, 432
Lychnis flos-cuculi, 172-
Lycopodium lucidulum, 1-
Lyellia aspera, 504

Macrocoma orthotrichoides, 145, 165; sullivantii, 145, 165
Macrodiplophyllum microdontum, 321-
Macromitrium amaniense, 227-; amboricum, *145*; amboroicum, 145; argutum, 145; atroviride, 145; bolivianum, *145*, *165*; brevihamatum, 145; cataractarum, 145; cirrosum, 145, 165; crassirameum, 145; crenulatum, 145; crispatulum, 145; didymodon, *145*, *165*; erectopatulum, *145*, *165*; filiforme, *145*, *165*; gigasporum, 145, 165; gigasporum var. brachypus, 145; glabratum, 145; herzogii, 145; hornschuchii, 145; incrustatifolium, 227-; liberum, *145*, *165*; longifolium, 145; macrosporum, *145*, *165*; macrothele, 145; nubigenum, 145; obtusum, *145*, *165*; pentastichum, 145; pinnulatum, 145; punctatum, 145, 165; refractifolium, 145; richardii, 145, 165; rusbyanum, 145, 168; solitarium, 145; solitarium var. brevipes, 145; stellulatum, 145; subbrevihamatum, 145; subcrenulatum, 145; subdiscretum, 145; sublaeve, 145; subscabrum, 145; swainsonii, 145; tocaremae, 145; tongense, 227-; tumidulum, 146; ulophyllum, 146; validum, 146; vesiculatum, 146, 168
MAGILL, R. E. Mosses of Big Bend National Park, Texas, 269
MAGILL, R. E. & R. KIRKPATRICK. Notes on Kansas Mosses, 101
Maine. distribution of Grimmia cribosa in, 356*, 357
Mandoniella spicatinervia, 155, 164
Manitoba. *See* Canada
Mannia fragrans, 432; pilosa, 432
MARAVOLO, N. C. *See* ALBERTINE, K. H. et al.
Marchantia polymorpha, 22, 186, 433
Marchantiaceae, 186, 433
Marchantiales, 432
Marsupella brevissima, 428; emarginata, 428; sparsifolia, 428; sphacelata, 428
Martin, W., 374*
Martinique. *See* West Indies
Mastodia tesselata, 396
MCFARLAND, K. D. & W. A. WISTENDAHL. Relationship of Terrestrial Mosses to Forested Flood Plains in Southeastern Ohio, 64
MCKNIGHT, B. N. *See* C. B. ARZENI & B. N. MCKNIGHT
Meesia triquetra, 444; uliginosa, 444
Meiothecium commutatum, 157; tenerum, 157
Merceya cataractae, *165*
Mesodon onustus, 157, 162
Metahygrobiella **tubulata**, comb. nov., 514
metal ions. effect on ascospore viability, 172
Meteoriaceae, 149
Meteoriopsis minuta, 149, 165; onusta, 149; patens, 149; patula, 149, 163; patula var. angustifolia, 149; pseudo-straminea, 149, 165; pseudo-straminea var. patula, 149, 165; recurvifolia, 149; reflecto-mucronata, 149, 165, 166; remotifolia, 149; remotifolia var. latifolia, 149; subrecurvifolia, 149
Meteorium ancistrodes, *247*; cladoniella, *149*, *165*; filiferum, *150*, *165*; fusco-viride, *149*, *165*; genuflexum, *247*; illecebrum, 149, 165, 166; lonchotrichum, *149*, *165*; minutum, *149*, *165*; perinflata, *150*, *165*; reflecto-mucronatum, *149*, *165*; stramineum, *149*, *165*; stramineum var. patulum, *149*, *165*
methylgyrophic acid, 224-
methylolivetoric acid, 42
Metzgeria conjugata, 1-, 432; crassipilis, 101; fruticulosa, 432; hamata, 432; pubescens, 432
Metzgeriaceae, 432
Metzeriales, 431
Metzleria longiseta, *129*, *165*; spiripes, *129*, *165*
Mexico. distribution of Ramalina leptocarpha and R. subleptocarpha in Baja California, 366*, 367*, 368, 369; distribution of Rozea in, 406-; new species of lichen from Nuevo Leon, 350; notes on Pottiaceae, 227; notes on tropical mosses in, 511; sensitivity of lichens [collected in Mexico]

to nitrogen dioxide, 103; Weissia rostellata var. phascoides new to, 384
Microdus exiguus, 131, 163; kunzeana, 131, 163; miquelianus, 495, 497*
microphyllinic acid, 42-
Microthamnium andicola, *159*, *165*; capillirameum, *159*, *165*; elegantulum, *159*, *165*; langsdorffii, *159*, *165*; oxystegum, *159*, *165*; reptans, *159*, *165*; thelistegum, *159*, *165*; viridicaule, *159*, *165*
microtopography. effects on establishment of a mixture of bryophytes from vegetative fragments, 199
Mielichhoferia, 122-, 480-; andina, 140; angustata, 140; aurifolia, *141*, *165*; boliviana, 140; brevicaulis, 140; bryocarpa, 140; campylocarpa, 140, 165; campylotheca, *140*, *165*; castanea, 140; clavitheca, 140; cygnicolla, 141; decurrens, 141; elegans, 141; emergens, 141; gracilis, 141; herzogii, 141; kerguelensis, *140*, *165*; lindigii, 141; lonchocarpa, 141, 165; longipes, 141; longiseta, 141; macrodonta, 141; macrospora, 141; microdonta, 141; micropoma, 141; mielichhoferi, 355-; minutifolia, 141; minutissima, 141; modesta, 141; modesta fo. viridis, 141; pohlioidea, 141; pusilla, 141; pusilla var. macrocarpa, 141; rigidula, 141; secundifolia, 141; seriata, 141; sericea, 141; spiniforme, *165*; subcampylocarpa, 141; subclavitheca, 141; subglobosa, 141; submacrodonta, 141; ventricosa, 141
MILLER, H. A. (review) K. W. ALLISON & J. CHILD, The Liverworts of New Zealand, 548
MILLER, N. G. & L. J. H. AMBROSE. Growth in Culture of Wind-Blown Bryophyte Gametophyte Fragments from Arctic Canada, 55
Mimosa spp., 269-
Missouri. new distributional records of bryophytes for, 100
Mittenothamnium andicola, 159, 165, 167; **burelae**, comb. nov., 159, 168; capillirameum, 159, 165, 167; decurrens, 159, 167, 168; diminutivum, 159, 165, 167, 168; elegantulum, 159, 165, 168; heterostachys, 159, 167; **hookerioides**, comb. nov., 159, 167; humile, 159, 168; langsdorffii, 159, 164, 165, 167; modestum, 159, 168; oxystegum, 159, 165; pendulinum, 159, 167; plumosum, 159, 167, 168; reptans, 159, 165, 167, 168; robustiusculum, 160, 167; simorhynchum, 160, 168; squarrosulum, 512; stigmopyxis, 160, 167, 168; subcampaniforme, 160, 168; subobscurum, 160, 167; **tenellum**, nom. nov., 160, 167; versipoma, 160, 168; viscidulum, 160, 167
Mniaceae, 142, 218
Mniobryum bolivianum, 141; wahlenbergii, 141, 162
Mnium. *See* also Plagiomnium & Rhizomnium; affine, 1-, 102, 241, *485*, 488-; arizonicum, 445; blyttii, 445, 484; ciliare, 95-; cuspidatum, 1-, 64-, 95-, 339-; hornum, 95-; ligulatum, *142*, *165*; lycopodioides, 504; marginatum, 1-, 484; orthorrhynchum, *484*; punctatum, 1-; punctatum var. elatum, 339-; riparium, 218; rostratum, 142; rostratum var. lingulatum, 142, 165; sp., 55-; spiniforme, *142*; spinosum, 504; spinulosum, 218; stellare, 95-; thompsonii, 445, 484; undulatum, 227-
Moenkemeyera obtusifolia, 128; termitarum, 128
Moerckia blytii, 186, 314-
Molendoa boliviana, *135*, *165*; boliviana var. brevifolia, *135*, *165*; cucullata, 135, 169; herzogii, *135*, *166*; sendtneriana, 135, 165, 166, 217; sendtneriana var. sendtneriana, 227; sendtneriana var. tenuinervis, 227
Montana. effects of pulp mill emissions on lichens in, 248; Sphagnum in, 508
morphogenesis. *See* growth
MOYLE, S. M. *See* BREIL, D. H. & S. M. MOYLE
MUELLER, D. M. J. Geocalyx graveolans, 268*
MUELLER, D. M. J. (review) J. D. HAYNES. Botany. An Introductory Survey of the Plant Kingdom, 264
MUELLER, D. M. J. Marchantia domingensis, 580*
MUELLER, D. M. J. & F. SEABURY. Funaria hygrometrica, 124*
Muhlenbergia spp., 269-
MURRAY, B. M. *See* HORTON, D. G. & B. M. MURRAY
Mycocalicium parietinum, 399; subtile, 399
Mylia anomala, 184, 428; taylori, 428
Myurella julacea, 55-, 321-, 445, 484; sibirica, 85, 106-, 321-, 504; tenerrima, 445, 484

NAIR, H. & V. RAGHAVAN. Cytokinin Induced Bud Formation in Microdus miquelianus, 495
Nardia, 428; geoscyphus, 184, 314-
Nardus stricta, 47-
NASH, T. H. Sensitivity of Lichens to Nitrogen Dioxide Fumigations, 103
NASH, T. H. *See also* THOMSON, J. W. & T. H. NASH

Nebraska. contributions toward a moss flora of, 241
Neckera cyathocarpa, *150*, *166*; disticha, *150*, *166*; eucarpa, 150; eucarpa var. secundifolia, 150; heteroclada, *151*, *166*; jamesonii, 150, 166; lindigii, 150; marchelii, 150; microtheca, *150*, *166*; trabeculata, 150; undulata, *151*, *166*
Neckeraceae, 150
Neckeropsis disticha, 150, 166, 511; microtheca, 150, 166; undulata, 151, 166
Neocardotia, 229; subnigra, 229
Neohypnella, 83-
Neosharpiella turgida, 138, 166
Nepal. distribution of Rozea in, 417
Nephroma arcticum, 453; expallidum, 453
Newfoundland. *See* Canada
New Hampshire. distribution of Grimmia cribosa in, 356*, 357
New Mexico. sensitivity of lichens [collected in New Mexico] to nitrogen dioxide, 103
News and Notes, 123, 266, 383
New York. distribution of Grimmia cribosa in, 356*, 357
New Zealand. Parmelia brattii in, 42
Nicaragua. Stenodictyon wrightii in, 83
Nicotiana glauca, 269-
nitrogen dioxide sensitivity of lichens to, 103
Nolina microcarpa, 269-
Normandina pulchella, 305
norstictic acid, 224-, 296-, 353, 385-, 500
North Carolina. distribution of Grimmia cribosa in, 356*, 357; Lophozia heterocolpa, new disjunct for, 222
Northwest Territories. *See* Canada
Nova Scotia. *See* Canada
Norwellia, 314-

Obituary. Yasuhiko Asahina, 258; William Martin, 374; Antero Vaarama, 377
Ochrobryum gardneri, 132, 166; gardnerianum, *132*, *166*; obtusifolium, 132; stenophyllum, 132
Ochrolechia frigida, 453; upsaliensis, 453
Octoblepharum albidum, 132; fragillimum, 132; pulvinatum, 132
Odontoschisma prostratum, 1-, 266-
Ohio. Hypotrachyna showmanii in, 78; relationship of terrestrial mosses to forested flood plains in southeastern, 64; seasonal growth of Parmelia caperata in southeastern, 360
Oligotrichum flacatum, 501-; hercynicum, 504
Oncophorus virens, 445, 484, 504; wahlenbergii, 445, 484
Ontario. *See* Canada
Opegrapha betulina, 399; bonplandi, 359; diaphora, 359; herbarum, 399; herpetica var. rufescens, *399*; lichenoides, 359; pulicaris, 359; rufescens var. rufescens, 399
Opuntia compressa, 1-; rufida, 269-
orcinol depsides, 43, 45*
Oregon. distribution of Ramalina subleptocarpha in, 367*, 368
Oreoweisia ampliata, 131; bogotensis, 131, 166; laxiretis, 131; ligularis, *131*, *166*; lechleri, 131; tunariensis, 131
Orphniospora atratra, *389*
Orthodontium confine, *141*, *166*; gracile, 141, 166, 168; longisetum, *141*, *166*; pellucens, 141, 166; setaceum, *141*, *166*
Orthostichidium excavatum, *149*, *166*; orthostichella, *150*, *166*; pentagonum, 149, 166
Orthostichopsis crinita, 149; rusbyana, 149, 166
Orthothecium, 406; acuminatum, 484; chryseum, 445, 484; diminutivum, 220; lapponicum, 484; lapponicum var. acuminatum, 484; rufescens, 484; sp., 55-; strictum, 220, 484
Orthotrichaceae, 145, 218, 290
Orthotrichum anomalum, 218, 445; apiculatum, *146*, *166*; cupulatum, 218; diaphanum, 86-, 290; elongatum, 146; emersulum, 146; epilosum, 146; exsertisetum, 146; jamesianum, 445; laevigatum fo. macounii, 445; liliputanum, 146; mandonii, 146; obtusifolium, 218, 445; pallens, 484; pariatum, 146; parvum, 146; patulum, 146; praeruptorum, 146; psychrophilum, 146; pulchellum, 445, 504; pumilum, 290; pycnophyllum, 290; pylaisii, 445; quenoae, 146; rupestre, 146; sordidulum, 146; speciosum, 208-, 484; speciosum var. elegans, 445; speciosum var. speciosum, 445; tacacomense, 146; undulatum, 146; verrucosum, 146; wagneri, 146, 166
Osorio, H. S. Contribution to the Lichen Flora of Argentina VIII. Lichens from Punta Lara, Buenos Aires Province, 358
Oxydendrum arboreum, 222-
Oxyrrhynchium aquaticum, *156*, *166*; clinocarpum, 155; clinocarpum var. brevisetum, 155; rugisetum, 155; scabripes, 156, 162
Oxystegus fallax, 135, 169; syrrhopodontoides, 135, 169
Østhagen, H. Nomenclatural Note on Cladonia nemoxyna, 242

Pachyospora mutabilis, 399; verrucosa, 399
Palamocladium leskeoides, 156, 166
Paludella squarrosa, 445
Pannaria pezizoides, 453; pityrea, 305
Paolillo, D. J. & M. Cukierski. Wall Developments and Coordinated Cytoplasmic

Changes in Spermatogenous Cells of Polytrichum (Musci), 466
Paolillo, D. J. *See* also French, J. C. & D. J. Paolillo
Papillaria appressa, *150*, *166*; brotheri, 150; cladomniella, *149*, *166*; deppei, 150; imponderosa, 150; imponderosa var. foveolata, 150; lonchotricha, *149*, *166*; nigrescens, 150, 166; squamatula, 150; tenella, 150
Paraleucobryum albicans, 131, 165
Parmelia sect. Cyclocheila, 506; alpicola, 453; aspera, 296-; bolliana, 225, 306; **brattii**, sp. nov., 42; caperata, 172-, 306, 360; caroliniana, 506; centrifuga, 453; cetrata, 359; chlorochroa, 401, 453; consors, 359; conspersa, 1-, 306; coronata, 306; crinita, 306; delisei, 42-; dierythra, 306; elegantula, 453; exasperatula, 453; flaventior, 306; formosana, 296-, 395; fraudans, 453; glomellifera, 42-; hababiana, 306; hypoleucites, 224-, 306; infumata, 453; kurokawae, 306; laevigata, 306; lindmanii, 359; lineola, 306; livida, 296-; **loxodella**, sp. nov., 43; loxodes, 42-; madagascariacea, 1-; margaritata, 306; microsticta, 359; nigropsoromifera, *225*; novomexicana, 307; perreticulata, 307; physodes, 172-; praesignis, 103-, 307; psoromifera, 307; pulla, 42-; pustulifera, 307; reticulata, 307, 359; rutidota, 307; saxatilis, 172-; spp., 172-; stygia, 453; subbalansae, 359; subolivacea, 307; subpraesignis, 225; subrudecta, 307; subtinctoria, 307; sulcata, 172-, 359, 453; taractica, 307, 453; tasmanica, 307; tubulosa, 172-; ulophyllodes, 307; verrucosa, 399; verruculifera, 42-; waiporiensis, 42-
Parmelina horrescens, 78-; spumosa, 78-
Parmeliopsis aleurites, 307; ambigua, 453
Parmotrema hypotropum, 224-; louisianae, 224
Paronychia argyrocoma, 222-
Peccania, 307
Pellia, 466-; endiviifolia, 186, 432; epiphylla, 1-, 22-, 432; neesiana, 432
Peltigera aphthosa var. aphthosa, 401, 453; aphthosa var. leucophlebia, 453; canina, 1-, 453; canina var. rufescens, 307; canina var. spuria, 307; elisabethae, 454; horizontalis, 454; leucophlebia, 401; malacea, 454; rufescens, 454; scabrosa, 454; spp., 172-; spuria, 454; venosa, 454
Peltula bolanderi, 308; euploca, 308; michoacanensis, 308; obscurans var. deserticola, 308; obscurans var. hassei, 308; obscurans var. obscurans, 308; omphaliza, 308; placodizans, 308; polyspora, 308; radicata, 308, 309*; richardsii, 309
Pennsylvania. Buellia dialyta in, 389; Hypotrachyna showmanii in, 80
perlatolic acid, 42-, 385-, 506
Pertusaria alpina, 454; dactylina, 454; pocillaria, *392*
Peru. Leptodontiella and Leptodontium spp. in, 20
Phacelia ranunculacea, 1-
Phaeocalicium curtisii, 402; populneum, 402
Phascum cuspidatum, 284
Philonotis, 122-; angulata, *144*, *166*; asperrima, 144; boliviana, *144*, *166*; boliviensis, 144; breviseta, *143*, *166*; calliantha, 144; clavicaulis, 144; curvata, 144, 166; eurydictyon, 144; filiramea, 144, 161; fontana, 290, 485; fontana var. fontana, 445; fontana var. pumila, 445; fontanella, 144, 166; gardneri, 144; glaucescens, 144, 162, 166; gracilenta, *144*, *166*; guyabayana, *144*, 161, *166*; hastata, 144, 166; laxissima, *144*, *166*; lignicola, 144; macrotheca, *143*, *166*; minutissima, *144*, *166*; muhlenbergii, 218; operta, 144; pagionifolia, *144*, *186*; parallela, 144; pellucidiretis, 144; pinnulata, *144*, *166*; pugionifolia, 144, 162; rufiflora, 144, 166; scabrifolia, 144, 162, 166; tenella, *144*, *166*; uncinata var. gracilenta, 144, 166
phosfon D (2,4-dichlorobenzyltributylphosphonium chloride). effects on plastid ultrastructure in Marchantia, 22
photosynthesis in lichens, 248
Phragmites communis, 269-
Phyllodoce empetriformis, 314-
Phyllogoniaceae, 150
Phyllogonium buchtienii, 150; fulgens, 150; viscosum, 150
Physcia adscendens, 454; aipolia, 310, 454; alba var. obsessa, 359; albinea, 310; caesia, 454; callosa, 310; convexa, 310; dubia, 86-, 310; elaeina, 310; grisea, 310; intermedia, 296-, 454; luganensis, 296-; mexicana, 310; nigricans, 310; orbicularis, 86-, 310; sciastra, 454; setosa, 310; stellaris, 310; syncolla, 310, 359; tenella, 86-; tribacoides, 311, 359; wainioi, 311, 454
Physcomitrium acuminatum, 101-; pyriforme, 102, 287, 515; turbinatum, 64-, 101-; turgidum, *138*, *166*
Physconia muscigina fo. alpina, 454; muscigina fo. muscigena, 454
Physiology. *See* also growth; effects of gibberellin and growth retardants on plastid ultrastructure, 22; influence of S-ethyl dipropylthiocarbamate on germinating Polytrichum spores, 332; lipid composition of mosses, 339; protein storage material in a moss, 188
phytogeography. additions to the bryophyte

flora of Colorado, 215; a new Cyathodium from India, 234; annotated checklist of hepatics of Idaho, 422; annotated checklist of mosses of Bolivia, 125; a Trichostomum from Texas, new to the United States, 80; contributions toward a moss flora of Nebraska, 241; distribution of Grimmia cribosa in North America, 355, 356*; distribution of Rhacomitrium lawtonae, 239*; Encalypta brevipes and E. mutica, gymnostomous species new to North America, 321, 328*; Hepaticae from west-central Alberta, Canada, 180; Hypotrachyna showmanii, a new lichen from eastern North America, 78; identity of Pterygoneurum arcticum with P. lamellatum, 221; Indusiella andersonii in northern Africa, 98; Lichenes Canadenses Exsiccati, 385; lichens from Punta Lara, Buenos Aires Province, Argentina, 358; Lophozia heterocolpa in North Carolina, 222; macrolichens of Big Bend National Park, Texas, 296; moss-lichen flora of St. Elias-Kluane Ranges, southwestern Yukon, Canada, 437; mosses new to the Yukon from the Ogilvie Mountains, Canada, 501; mosses of Big Bend National Park, Texas, 269; neotropical Polytrichaceae, 93, 521; new and additional lichen records for Texas, 224; new distributional records of bryophytes for Arkansas and Missouri, 100; new or additional moss records from Newfoundland, Canada, 84; new or additional moss records of northern Ellesmere Island, Arctic Canada, 480; new species and new combinations of hepatics from Chile, 514; notes on Kansas mosses, 101; notes on Pottiaceae in Middle America, 227; notes on tropical mosses of Mexico, 511; Pyxine azorea is Parmelia caroliniana, 506; Ramalina farinacea in Hawaii, 499; Ramalina leptocarpha and R. subleptocarpha, a fertile-sorediate species pair, 364, 366*, 367*; revision of the genus Rozea (Musci), 406; Schistidium holmenianum, sp. nov. from Arctic North America, 208; Schofieldia, a new hepatic from the Pacific Northwest, 314; Seligeria brevifolia, a moss new to North America, 106; Seligeria careyana, a new species from the Queen Charlotte Islands, western Canada, 231; some mosses from Kuwait, 515; Sphagnum in the northern Rocky Mountains of the United States, 508; Sphagnum pylaesii in Greenland, 47; Syrrhopodon annotinus (Musci; Calymperaceae), a new species from Brazil, 518; three new lichens from the Southwest, 350; Weissia rostellata var. phascoides new to Mexico, 384

Picea glauca, 437-; mariana, 437-

Pilopogon gracilis, 131; gracilis var. comigera, 131; gracilis var. divaricatus, 131; gracilis var. parvus, 131, 166; holomitrius, 131; liliputanus, *129*, *166*; longifimbriatus, *131*, *166*; nanus, 131; richardii, *130*, *166*; tiquipayae, 131

Pilosium chlorophyllum, 157, 168

Pilotrichella angustifolia, 150; cyathipoma, 150; cyathipoma var. laxiretis, 150; dimorpha, *149*, *166*; flexilis, 150, 166; flexilis var. minor, 150; perinflata, *150*, *166*; reflecto-mucronata, *149*, *166*; turgescens, *150*, *166*; versicolor, 150; viridis, 150

Pilotrichum fendleri, 512; genuflexum, *247*

Pinnatella ochracea, 151

Pinus cembroides, 269-; edulis, 215-; ponderosa, 269-; spp., 172-; strobus, 78-

Pirea pohlii, *149*, *166*

Pireella pohlii, 149, 166

Plagiobryum demissum, 321-, 445, 504; zierii, 321-, 445, 504

Plagiochila asplenioides, 1-, 185, 430; **cymbiformis**, sp. nov., 514; major, 430; rhizophora, 430; satoi, 430; semidecurrens, 430; sharpii, 1-

Plagiochilaceae, 185, 430

Plagiomnium ciliare, 485; ellipticum, 445, 485; medium, 485; medium ssp. curvatulum, 485; rugicum, *485*

Plagiopus oederi, 144; oederiana, 445, 485

Plagiotheciaceae, 157, 220

Plagiothecium andicola, *157*, *166*; andinum, 157, 166; bolivianum, *157*, *166*; cavifolium, 220; conostegium, 157; denticulatum, 95-; fallax, *157*, *166*; herzogii, 157, 166; microsphaerothecium, 157; mollicaule, 157; novogranatense, 157; roeseanum, *220*; submollicaule, 157

Plantago major, 172-

plastids. effects of gibberellin and growth retardants on ultrastructure in Marchantia, 22

Platanus occidentalis, 64-

Platydictya confervoides, 1-; jungermannioides, 85, 106-, 485, 504

Platygyrium, 406-; fuscoluteum, 294; repens, 95-

Platyhypnidium aquaticum, 156, 166, 167

Platyneurum, 122-

Plectocolea crenuliformis, 1-

Pleuridium andinum, 128

Pleurochaete squarrosa, 284

Pleuroclada albescens, 186, 314-

Pleuropus bonplandii, *156*, *166*

Pleurozium, 406-; schreberi, 157, 164, 199-, 222-, 445

Poa glauca, 437-

Pogonatum abbreviatum, 160; arcuatum, 160;

capillare, *485*; cylindrotheca, 160; dentatum, 485, 504; distantifolium, 160; germainii, 160, 166; glabrum, 521; laxirete, 160; nanum, 93-; oligodus, 160; plurisetum, 160, 166, 169; polycarpum, *160, 166*; polycarpum var. angustatum, *160, 166*; purpurascens, 160; semiangulatum, *521*; subbifarium, 160; tortile, 160; urnigerum, 445, 504

Pohlia. *See* also Anomobryum & Mniobryum; 122-; annotina, 445; annotina var. decipiens, 218; apolensis, 141, 169; atropurpurea, 445, 504; bulbifera, 218, 445; **clavicaulis**, comb. nov., 141, 169; cruda, 141, 169, 445, 485; drummondii, 218, 289; **loriformis**, comb. nov., 141, 169; mandonii, 141; nutans, 22-, 55-, 446; papillosa, 141, 169; pluriseta, 141; proligera, 446; rothii, 446; rusbyana, 141, 162; schistocola, 141, 162; spectabilis, 142, 169; **subleptopoda**, comb. nov., 142, 169; verrucosa, 142, 162; wahlenbergii, 446, 485

Polianthes virginica, 1-

Polymerodon andinus, 131

Polypodium virginianum, 222-

Polytrichaceae, 93, 98-, 160, 521

Polytrichadelphus aristatus, 160; bolivianus, 160; croceus, *521*; cuspidirostrum, 160, 166; ericoides, *93*; grossidens, 160, 162; integrifolius, 160, 162; juniperaceus, *521*; magellanicus, 521; mitis, 160; **pseudopolytrichum**, comb. nov., 521; rubiginosus, 161; semiangulatus, *521*; trianae, 161; umbrosus, 161

Polytrichastrum alpinum, 446; tenellum, 161, 166

Polytrichum. *See also* Polytrichastrum; 122-, 457-, 466-; algidum, 504; aloides, 93-; angusticaule, *161, 166*; antillarum, *161, 166*; aristiflorum, *161, 166*; commune, 1-, 93-, 95-, 161, 188-, 199-, 201*, 242, 332, 335*, 339-, 446, 488, 490*; commune var. jensenii, 76; cuspidigerum, *161, 166*; cuspidirostrum, *166*; ericoides, 93, 94*; formosum, 95-, 222-; germainii, *160, 166*; hyperboreum, 76, 504; intermedium, *161, 166*; juniperinum, 98-, 161, 166, 242, 339-, 446, 468*, 470*, 472*, 474*, 476*, 478*; juniperinum var. tumescens, 161; magellanicum, *521*; nanum, 93-; ohioense, 95-; patens, *161, 166*; piliferum, 446, 485, 504; polycarpum, *166*; ramosum, 161; rhapidostegium, 521; secundulum, *161, 166*; secundulum var. angusticaule, *161, 166*; sexangulare, 446, 504; spp., 172-; strictum, 446; subnudum, 521; tenellum, *161, 166*

Populus, 241-; balsamifera, 437-; deltoides, 1-, 64-; palmeri, 269-; tremuloides, 437-

Porella cordaeana, 186, 431; navicularis, 431; pinnata, 95-; platyphylla, 1-, 95-, 431; platyphylloidea, 431; roellii, 431

Porellaceae, 186, 431

Porina chlorotica, 402; nucula, 359

Porothamnium comosum, 151; explanatum, 151, 166; fasciculatum, 151, 166, 169; gymnopodum, 151; neckeraeforme, 151, 166; ramosissimum, 151, 169; subexplanatum, 151; subramosissimum, 151

Porotrichodendron bolivianum, *151, 166*; gracile, 151; robustum, 151; superbum, 151, 166, 169

Porotrichopsis flacca, 151

Porotrichum amboroicum, 151; bolivianum, 151; explanatum, *151, 166*; fasciculatum, *151, 166*; longirostre, 151, 169; lorentzii, *151, 166*; macropoma, 151; microthecium, 151; pinnatelloides, 151; plicatulum, 513; sp., 512; strictum, 151; superbum, *151, 166*

Potamium longisetum, *158, 166*

Pottia arizonica, 284; arizonica var. mucronulata, 269-; davalliana, 516; lanceolata, 516; starckeana, 516

Pottiaceae, 98-, 132, 216, 279

Pouteria salicifolia, 358-

Prasiola borealis, 397, 398*; meridionalis, 398*, 399; sp., 398*; tesselata, 397

Preissia quadrata, 186, 433

Prionodon bolivianus, 148; cavifolius, *149, 166*; ciliatoserratus, *148, 167*; contortus, 148; densus, 148, 167; divaricatus, *148, 167*; filifolius, 148; flagellaris, *148, 167*; fusco-lutescens, 148, 167; laeviusculus, *148, 167*; luteo-virens, 148, 167; lycopodioides, 149, 166; patensissimus, *149*; patentissimus, 149; pendulus, *148, 167*; pinnatus, 149; ptychomnioides, 149; splendens, *149, 167*; tenerrimus, 149; undulatus, *148, 167*

Prionodontaceae, 148

propagation Sphagnum pylaesii, 52

Prosopis spp., 269

protein storage material in Dicranum, 188

Protobastenia rupestris, 454

Pseudevernia intensa, 311

Pseudocrossidium, 98-; pachygastrellum, 135, 161

Pseudocryphaea flagellifera, 148

Pseudodimerodontium bolivianum, 153, 165, 167, 168; minuta, 153, 168

Pseudoleskea amblystegiella, *153, 167*; andina, 154, 167; catenularia, *153, 167*; catenulatula, *154, 167*; minuata, *153, 167*; radicosa var. compacta, 446; rusbyana, *154, 167*

Pseudoleskeella nervosa var. sibirica, 218, 505; papillosa, 505

Pseudoparmelia, *506*; cryptochlorophaea, 224; caroliniana, *506*
Pseudotsuga menziesii, 269-
Psilopilum, 122-; aequinoctiale, 161, 162; antarcticum, 161; cavifolium, 321-; gymnostomulum, 161, 162; trichodon, 161
Psoroma hypnorum, 454
psoromic acid, 224, 296-
Ptelea, 215-
Pteridium, 466-
Pterigynandrum, 406-; filiforme, 505
Pterobryaceae, 149
Pterobryon climacioides, 149; densum, 149, 167
Pterobryopsis stolonacea, 149
Pterogonidium pulchellum, 158, 167
Pterogoniella pulchella, *158*, *167*
Pterygoneurum arcticum, *221*, 485; lamellatum, 221; ovatum, 446, 516; subsessile, 221-, 284, 446
Pterygynandrum filiforme, 85
Ptilidiaceae, 182, 424
Ptilidium californicum, 424; ciliare, 182, 425; pulcherrimum, 182, 425
Ptilium crista-castrensis, 446
Ptychomitriaceae, 144, 290
Ptychomitrium angustifolium, 144; chimborazense, 144; cochabambae, 144, 164; ferrugineum, 145, 164; incurvum, 222-, 290; leibergii, 290; mandonii, 145; papillosum, 145, 164; sellowianum, 145, 164
Pungentella levieri, *158*, *167*
PURSELL, R. A. On the Typification of Certain Taxa and Structural Variation Within the Fissidens bryoides Complex in Eastern North America, 35
Purshia, 215-
PYATT, F. B. Lichen Ecology of Metal Spoil Tips: Effects of Metal Ions on Ascospore Viability, 172
Pylaisia boliviana, 160; panduraefolia, 160; subjulacea, *411*
Pylaisiella, 406-; polyantha, 446; selwynii, 242
Pylaisiobryum, 406-
Pyxine azorea, *506*

Quebec. *See* Canada
Queen Charlotte Islands. *See* Canada
Quercus, 241-; emoryi, 269-; gravesii, 269-; grisea, 269-; lobata, 364-; macrocarpa, 241-; montana, 78-; petraea, 172-; rubra, 222-; spp., 172-

Racomitrium. *See* Rhacomitrium
Radula auriculata, 431; bolanderi, 431; complanata, 1-, 186, 431; obconica, 1-; obtusiloba, 431
Radulaceae, 186, 431
RAGHAVAN, V. *See* NAIR, H. & V. RAGHAVAN
Ramalina, 350-; calicaris, 364-; canaliculata, 364-; complanata, 311, 359; ecklonii, 311, 359, 364-; ecklonii fo. lobuligera, 359; farinacea, 364-, 365*, 499; irregularis, *368*; leptocarpha, 364-, 365*; linearis, 364-; menziesii, 368, *369*; F. soderiata, *368*; peruviana, 359; pollinaria, 311; prolifera, 359; reagens, 499-; reticulata, 368-; roesleri, 454; sinensis, 311; spp., 172-; subfarinacea var. reagens, 499-; **subleptocarpha**, sp. nov., 365*, 368
Ranunculus spp., 172-
RASTORFER, J. R. Caloric Values of Three Alaskan-Arctic Mosses, 76
RASTORFER, J. R. (review) S. W. GREENE, B. G. BELL, G. C. S. CLARKE & M. E. NEWTON, A Synoptic Flora of South Georgian Mosses, 122
Rauia bornii, *154*, *167*
Rauiella **bornii**, comb. nov., 154, 167; firmula, 154, 169; plumaria, 154, 165
Reboulia hemisphaerica, 1-, 186, 222-, 269-
Rechingeria cribellifera, 311
REDFEARN, P. L. A Trichostomum from Texas New to the United States, 80
REDFEARN, P. L. *See also* P. E. THOMAS & P. L. REDFEARN
REESE, W. D. & D. GRIFFIN. Syrrhopodon annotinus (Musci; Calymperaceae), a New Species from Amazonas, Brazil, 518
REESE, W. D. *See also* DELGADILLO M., C. & W. D. REESE
regeneration. *See* growth
reviews, 122, 264, 380, 547
Rhabdoweisia crispata, 131, 167; denticulata, *131*, *167*; fugax, 131, 167; lindigiana, *131*, *167*
Rhacocarpus australis, *147*, *167*; chlorotus, 147; excisus, 147; humboldtii, *167*; purpurascens, 147, 164, 167; squamosus, 148
Rhacomitrium, 122-; aciculare, 217; austrosudeticum, 137; brachypus, 137, 164; canescens, 485; canescens var. ericoides, 446; crispipilum, *137*, *167*; crispulum, 137, 164, 167; dimorphum, 137, 164; heterostichum, 217, 238-; heterostichum var. sudeticum, 446; lanuginosum, 446, 485; lawtonae, 238, 240*, distribution of, 239*; sublanuginosum, 138; sudeticum, 208-; varium, 238-
Rhacopilaceae, 147
Rhacopilum floridae, 147; intermedium, 147; tomentosum, 147; tomentosum var. papillosum, 147
Rhamphidium levieri, *128*, *167*; pygmaeolum, 135, 161
Rhaphidorrhynchium andinum, 158, 167, 168; capillifolium, 158, 167; decumbens, 158,

167; geheebii, 158; levieri, 158; lindigii, 158, 167; lindigii var. propaguliferum, *158*; oblique-rostratum, 158, 168; prominulum, 158, 167, 168; rueckeri, 158, 167
Rhaphidostegium andinum, *158*, *167*; aureoviride, *158*, *167*; caespitosum, *158*, *167*; capillifolium, *158*, *167*; chiquitanum, 158; chlorocormum, 158, 161; circinale, *158*, *167*; cuspidiferum, *158*, *167*; decumbens, *158*, *167*; densirete, 158; eurycystis, 158; flavidum var. breviapiculatum, *158*, *167*; galipense, *158*, *167*; geheebii, *158*, *167*; guarayum, 158; kegelianum, *158*, *167*; lamellosum, 158; lindigii, *158*, 167; lindigii var. propaguliferum, *158*, *167*; loxense, *158*, *167*; orthocarpum, *158*, *167*; proligerum, *157*, *167*; prominulum, *158*, *167*; rueckeri, *158*, *167*; tenerifolium, *158*, *167*; turgidulum, 158; undulatum, 158
Rhexophyllum, 229; laciniatum, 135, *229*; subnigrum, 229, 284
Rhizocarpon, 389; disporum, 454; geographicum, 454; macrosporum, 454; sphaerosporum, 454
Rhizogoniaceae, 142
Rhizogonium bolivianum, 142; spiniforme, 142, 165
Rhizohypnum acrorrhizon, *159*, *167*; andicola, *159*, *167*; breviusculum, *160*, *167*; capillirameum, *159*, *167*; decurrens, *159*, *167*; delicatulum, *160*, *167*; heterostachys, *159*, *167*; hookerioides, *159*, *167*; pelichucense, *155*, *167*; pendulinum, *159*, *167*; plumosum, *159*, *167*; reptans, *159*, *167*; robustiusculum, *160*, *167*; stigmopyxis, *160*, *167*; thelistegum, *159*, *167*; viridicaule, *159*, *167*; viscidulum, *160*, *167*
Rhizomnium gracile, 446
Rhodobryum beyrichianum, 142; caulifolium, *139*, *167*; grandifolium, 142, 162; lindigianum, *140*, *167*; roseum, 95-, 142. 289; verticillatulum, 142
Rhododendron catawbiense, 222-
Rhus typhina, 385-
Rhynchostegiella semitorta, *155*, *167*; toncolensis, *155*, *167*
Rhynchostegiopsis complanata, 153
Rhynchostegium albo-viridum, 156; ambiguum, 156; aquaticum, *156*, *167*; bolivianum, 156; brevirete, 156; callistomum, *156*, *167*; conchophyllum, 156; hirtipes, *155*, *167*; lamasicum, 156; limosum, 156; mandonii, 156; minutum, 156; planifolium, 156; planifolium var. tenue, 156; scariosum, 156; serrulatum, 95-, 156, 167; subrusciforme, 156; tocaremae, 156
Rhytidiaceae, 160
Rhytidium rugosum, 160, 446
Riccardia latifrons, 186, 432; multifida, 432; palmata, 101, 432
Riccia austinii, 1-, 220; beyrichiana, 1-, 433; crystallina, 433; frostii, 433; hirta, 1-
Ricciaceae, 220
Ricciocarpus natans, 433
Riella, 466-
Rigodium leptodendron, 151
Rinodina mniaraea, 454; nimbosa, 455; roscida, 455; turfacea, 455
Rozea, 409; andrieuxii, *413*; bourgaeana, *415*; chrysea var. bourgaeana, 414*, 415; chrysea var. chrysea, 411; chrysea var. chrysea fo. chrysea, 411, 412*; chrysea var. chrysea fo. **andrieuxii**, stat. nov., 412*, 413; diversifolia, 409; fulva, 409, *419*; **kenoyeri**, sp. nov., 418*, 420; microcarpa, 409, 419; microcarpa fo. **elongata**, fo. nov., 418*, 420; microcarpa fo. microcarpa, 418*, 419; myura, *417*, 418*; petrophila, *409*; pterogonioides, 417, 418*; roseorum, 409; schimperi, 414*, *415*; stricta, *411*, 412*; subjulacea, 414*, 416; viridis, 414*, *415*
Rubus fruticosus, 172-
Rumex spp., 172-
Rundel, P. W. & P. A. Bowler. Ramalina leptocarpha and R. subleptocarpha: A Fertile-Sorediate Species Pair, 364

Saadawi, W. S. Some Mosses of Kuwait, 515
Saarinen, R. *See* Valanne, N. et al.
Saelania glaucescens, 86, 446, 485
salazinic acid, 296-, 500
Salix, 358-; glauca, 437-; nigra, 64-; planifolia, 437-; polaris, 437-; reticulata, 437-; scouleriana, 437-; spp., 269-
Salvia regla, 269-
Sanderson, C. *See* Sheridan, R. P. et al.
Sanicula canadensis, 64-
Sapium, 358-
Saskatchewan. *See* Canada
Saxifraga michauxii, 222-; oppositifolia, 55-
scanning electron micrographs. peristome of Ctenidiadelphus cylindricarpus, 408*; peristome of Entodon macropodus, 410*; peristome of Hypnum fertile, 408*; peristome of Rozea subjulacea, 408*; spores of Encalypta brevipes & E. mutica, 327*
Scapania americana, 429; apiculata, 184, 429; bolanderi, 429; curta, 184, 428; cuspiduligera, 429; degenii var. dubia, 185; dentata, 429; evansii, 429; glaucocephala, 185, 429; gymnostomophila, 435; irrigua, 185, 428; mucronata, 185, 428; nemorosa, 1-, 266-, 428; paludosa, 185, 428; scandica, 429; subalpina, 428; uliginosa, 429;

umbrosa, 428; undulata, 1-, 185, 428; undulata var. oakesii, 429
Scapaniaceae, 184, 428
Schistidium alpicola, 208-, 485; andinum, *136*, *168*; angustifolium, *137*, *168*; apocarpum, *137*, *168*, 485; apocarpum var. gracilis, 208-; apocarpum var. nigrescens, 208-; boreale, 208-; chocayae, *137*, *168*; fontanum, *137*, *168*; gracile, 485; **holmenianum**, sp. nov., 208, 209*, 210*, 211*, 485; malacophyllum, *137*, *168*; praemorsum, *137*, *168*; streptophyllum, *137*, *168*; subpraemorsum, *137*, *168*; tenerum, 485; tunariense, *137*, *168*
Schlenk, H. *See* Swanson, E. S. et al.
Schlotheimia angustata, 146; argentinica, 146; breviseta, 146; campylopus, 146; fusco-viridis, 146; jamesonii, *146*, *168*; krauseana, *146*, *168*; krausei, 146, 168; lindmanii, 146; longicaulis, 146; pilomitria, 146; rugifolia, 146, 168; rusbyana, 146; sprengelii, *146*, *168*; sublevifolia, 146; torquata, 146, 168; trichomitria, 146; vesiculata, *146*, *168*
Schofield, W. B. *See* Vitt, D. H. & W. B. Schofield
Schofieldia, gen. nov., 314; **monticola**, sp. nov., 315, 316*, 317*
Schraderobryum ulicinum, 158, 168
Schroeterella zygodonta, 158
Schwetschkea boliviana, *153*, *168*; minuta, *153*, *168*
Sciaromium crassinervatum, 155; holoneuron, 155
Scirpus caespitosus, 47-
Scopelophila cataractae, 135, 165; ligulata, 355-
Scorpidium scorpioides, 155, 446; turfaceum, 155, 164; turgescens, 208-, 446, 485
Scouleria aquatica, 208-
Scutia, 358-; buxifolia, 358-
Seabury, F. *See* Mueller, D. M. J. & F. Seabury
Sedum spp., 172-
Seligeria acutifolia, 231-; brevifolia, 106; campylopoda, 86, 231-, 505; cardotii, 231-; **careyana**, sp. nov., 232, 233*; diversifolia, 505; donniana, 101, 505; oelandica, 505; polaris, 231-, 485, 505; pusilla, 231-, 484; recurvata, 86, 106-; subimmersa, 446, 505; tristichoides, 505
SEM. *See* scanning electron micrographs
Sematophyllaceae, 157
Sematophyllum andinum, *158*, *168*; aureoviride, 158, 167; brachyacrum, 158, 161; caespitosum, 158, 167, 168; carolinianum, 1-; chrysostegum, 158; circinale, *158*, *168*; cucullatifolium, *158*, *168*; cuspidiferum, 158, 167; demissum, 95-; flavidum, 158, 167; galipense, 158, 167; longisetum, 158, 166; loxense, *158*, *168*; oblique-rostratum, *158*, *168*; orthocarpum, 158, 167; prominulum, *158*, *168*; pungens, *157*, *168*; sp., 158, 168; subsimplex, 158; tenerifolium, 158, 167; tenuicarpum, 158; ulicinum, *158*, *168*
S-ethyl dipropylthiocarbamate. Effect on chlorophyll and carotenoid production and chloroplast ultrastructure of germinating Polytrichum spores, 332
Sharp, A. J. *See* Juárez G., L. G. & A. J. Sharp
Sheard, J. W. (review) A. L. Smith, Lichens, 382
Sheridan, R. P., C. Sanderson, & R. Kerr. Effects of Pulp Mill Emissions on Lichens in the Missoula Valley, Montana, 248
Showman, R. E. Seasonal Growth of Parmelia caperata, 360
Shuster, R. M. (review) S. R. Gradstein, A Taxonomic Monograph of the Genus Acrolejeunea (Hepaticae), 380
Sigmatella stigmopyxis, *160*, *168*
Silene maritima, 172-
Simplicidens andicola, 128; plumula, 128
Singh, D. K. *See* Udar, R. & D. K. Singh
Siphula ceratites, 402
Smith, A. L. Lichens (review), 382
Smith, C. W. *See* Bowler, P. A. & C. W. Smith
Smith, G. L. Neotropical Polytrichaceae IV, 93
Smith, G. L. Neotropical Polytrichaceae V, 521
Solenostoma atrovirens, 427; cordifolium, 184, 427; hyalinum, 428; obovatum, 428; pumilum, 427; pumilum subsp. polaris, 427; rubrum, 427; sp., 55-; sphaerocarpum, 427; triste, 184, 428
Solorina crocea, 455; octospora, 455; saccata, 455; spongiosa, 455
Sorbus spp., 172-
Southbyaceae, 185
South Georgia, mosses of (review), 122
Søchting, U. *See* Johnson, I. & U. Søchting
Sparganium hyperboreum, 47-
spermatogenous cells of Polytrichum, 466
Sphaerocarpaceae, 432
Sphaerocarpus hians, 432; melanocarpus, 402
Sphagnaceae, 127, 215
Sphagnum acutifolium, *127*, *168*; americanum, *127*, *168*; angustifolium, 509; balticum, 127, 168; boliviae, 127; boliviae var. virescens, 127; capillaceum, 95-, 102; capillaceum var. tennellum, 508-; capillifolium, 127, 168; centrale, 509; compactum, 47-, 505; contortum, 509; erythrocalyx, *127*, 168; fallax, 508-; fimbriatum,

215, 509; fuscum, 215, 391, 446, 509; girgensohnii, 47-, 446, 505, 509; gracile, *127*, *168*; jensenii, 509; lenense, 505; lindbergii, 47-, 505; longifolium, *127*, *168*; magellanicum, 127, 168, 509; magellanicum fo. purpurascens, 127, 168; magellanicum var. molle, 127, 168; magellanicum var. pallescens, 127, 168; medium, *127*, *168*; medium var. pallescens, *168*; medium var. pallescens fo. densa, *127*, *168*; medium var. purpurascens, *127*, *168*; medium var. virescens, *127*, *168*; mendocinum, 509; meridense, 127, 168, var. patulum, 127; molle, 508-; molle var. limbatum, 127, 168; nemoreum, 505, 509; ovatum, *127*, *168*; palustre, 95-; papillosum, 47-, 199-, 201*; perichaetiale, 127, 168; peruvianum, *127*, *168*; platyphylloides, 127; platyphyllum, 47-, 50*, 509; plumulosum, 508-; pulchricoma, *127*, *168*; pylaesii, 47, 49*-51*, photographic survey of Greenland localities, 48*, "ramosum" type, 47-, "sedoides" type, 47-; recurvum, 127, 168, 508-; recurvum var. mucronatum, *127*, *168*; riparium, 505, 509; rubellum, 508-; russowii, 505, 509; sancto-josephense, 127; sedoides, 47-; sparsum, 127; squarrosum, 509; subcuspidatum, *127*, *168*; subsecundum, 95-, 509; tenellum, 47-; teres, 509; warnstorfii, 446, 510; wolfianum, 508-

Spiloma vitiligo, *403*

Spinacia oleracea, 332-

Splachnaceae, 138, 287

Splachnobryum crenulatum, 227-; obtusum, 227-, 287

Splachnum, 227-; ampullaceum, 22-; luteum var. luteum, 446; luteum var. melanocaulon, 447; sphaericum, 447

Sporastatia testudinea, 455

Squamarina lentigera, 455

Squamidium diversifolium, 150; filiferum, 150, 165; leucotrichum, 150; lorentzii, 150; macrocarpum, 150; nigricans, 150, 166; perinflatum, 150, 165, 166; turgidulum, 150

Stableria tennellia, *141*, *168*

Staphylea trifolia, 1-

STEERE, W. C. Identity of Pterygoneurum arcticum with P. lamellatum, 221

STEERE, W. C. & G. R. BRASSARD. Schistidium holmenianum, sp. nov. from Arctic North America, 208

Stegonia latifolia, 58-, 321-, 486; latifolia var. latifolia, 447, 505; latifolia var. pilifera, 447, 486

Stenodictyon nitidum, 83-; saxicola, 153; sericeum, *84*; **wrightii**, comb. nov., 83

stenosporic acid, 506

Stereocaulon alpinum, 455; glareosum, 455; grande, 455; rivulorum, 455; tomentosum, 455

Stereodon spiripes, *159*, *168*

Stereohypnum burelae, *159*, *168*; decurrens, *159*, *168*; elegantulum, *159*, *168*; humile, *159*, *168*; modestum, *159*, *168*; plumosum, *159*, *168*; reptans, *159*, *168*; simorhynchum, *160*, *168*; stigmopyxis, *160*, *168*; subcampaniforme, *160*, *168*; thelistegum, *159*, *168*; versipoma, *160*, *168*

Stereophyllum aptychopsis, 157; brevipes, 157; cultelliforme, 157; cupreum, 157; flaccisetum, *157*, *168*; fulvum, *419*; homalioides, 157; leptostegium, 157; leucostegum, 157; lindmanii, 157; myurum, 157; pseudo-radiculosum, *157*, *168*; subchlorophyllum, *157*, *168*

stictic acid, 296-

St. Kitts. *See* West Indies

Stokesiella praelonga, 156, 163

STOTLER, R. E. Saxicolous Bryophyte and Macrolichen Associations in Southern Illinois I. Little Grand Canyon, Jackson County, 1

Streptopogon bolivianus, *135*, *168*; calymperes, 135, 168; clavipes, 135; erythrodontus, 135, 168; heterophyllus, 135; lindigii, 135, 168; peruvianus, 135; rigidus, *135*, *168*; setiferus, *135*, *168*; spathulatus, 135

Streptotrichum ramicola, 16-, 135

Strigula elegans, 359

structural variation. in Fissidens bryoides complex, 37; in Sphagnum pylaesii, 51

substrate type. effects on establishment of a mixture of bryophytes from vegetative fragments, 199

sulfur dioxide, effect on lichens, 248

SWANSON, E. S., W. H. ANDERSON, J. L. GELLERMAN & H. SCHLENK. Ultrastructure and Lipid Composition of Mosses, 339

SWINSCOW, T. D. V. *See* KROG, H. & T. D. V. SWINSCOW

Symblepharis boliviana, *131*, *168*; lindigii, 131, 168

Synalissa symphorea, 311

Syringothecium nemodontium, 158

Syrrhopodon **annotinus**, sp. nov., 518, 519*; anomalus, *132*, *168*; argenticus, 132; brachystelioides, *132*, *168*; ciliolatus, *132*, *168*; circinatus, 132; densifolius, 132, 520; elatior, 132; gaudichaudii, 132; goyazensis, 132; hobsonii var. spinulosus, 132; incompletus, 132, 168, 512; leprieurii, 132; ligulatus, 132, 168; lycopodioides, 132, 513; macrophyllus, 132; martinii, 132; miquelianus, 132; ochroleucus, 132, 520; papillosus, 132; parasiticus, 520; prolifer, 132, 168; pusillus, 132; quintasii, 132,

168; rupicola, 520; scaber, *132*, *168*; scaberrimus, 132; serpentinus, 132; spininervis, 132; subdecolorans, 132; submacrophyllus, 132; tricolor, *20*, *134*, *168*; ulei, 520
systematics. a new combination for Cladonia stellaris, 363; a new Cyathodium from India, 234; a Trichostomum from Texas new to the United States, 80; Barbula bescherellei, etc., 215; correct name for the moss Duseniella, 246; Encalypta brevipes and E. mutica, 321; Fissidens bryoides complex in eastern North America, 35; Grimmia cribosa in North America, 355; Hypotrachyna showmanii, a new lichen from eastern North America, 78; identity of Pterygoneurum arcticum with P. lamellatum, 221; Indusiella andersonii, sp. nov. in Northern Africa, 98; Leptodontiella, gen. nov. and Leptodontium from Peru, 16; Lichenes Canadenses Exsiccati, 395; macrolichens of Big Bend National Park, Texas, 296; mosses of Big Bend National Park, Texas, 269; mosses of northern Ellesmere Island, Arctic America, 480; Neotropical Polytrichaceae, 93, 521; new and additional lichen records from Texas, 224; new species and new combinations of hepatics from Chile, 514; nomenclatural notes on Cladonia nemoxyna, 242; notes on Pottiaceae in Middle America, 227; Pyxine azorea and Parmelia caroliniana, 506; Ramalina leptocarpha and R. subleptocarpha, 364; recopilación de los musgos de Bolivia, 125; relationships of Hookeria wrightii and Stenodictyon sericeum, 83; revision of the genus Rozea, 406; Rhacomitrium lawtonae, 238; Schistidium holmenianum, sp. nov. from Arctic North America, 208; Schofieldia, a new hepatic from the Pacific Northwest, 314; Seligeria careyana, a new species from the Queen Charlotte Islands, western Canada, 231; Sphagnum pylaesii in Greenland, 47; Syrrhopodon annotinus, a new species from Brazil, 518; three new lichens from the southwest, 350

Tamariscella tripinnata, *154*, *168*
Tamarix gallica, 269-
Taraxacum spp., 172-
Tasmania. *See* Australia
Taxicaulis cylindraceus, *157*, *168*; stigmocarpus, *157*, *168*; subcylindraceus, *158*, *168*
Taxiphyllum deplanatum, 95-, 242; scalpellifolium, 512; taxirameum, 222-
taxonomy. *See* systematics
Tayloria, 227-; altorum, 138; cochabambae, 138; cochambae, *138*; froelichiana, 505; jamesonii, 138, 162; lingulata, 447, 505; mandonii, 138, 162; mnioides var. cavifolius, 447; mnioides var. mnioides, 447; moritziana, 139; scabriseta, 139
Teichodontium rusbyanum, *145*, *168*
Telaranea (subg. Neolepidozia) **oligophylla**, comb. nov., 514; (subg. Neolepidozia) **seriatitexta**, comb. nov., 514
Teloschistes chrysophthalmus, 311, 360; flavicans, 311
Tennessee. distribution of Grimmia cribosa, 356*, 357
Teretidens flaccidus, *131*, *169*
Terminalia australis, 358-
Tetraphis pellucida, 1-
Tetraplodon angustatus, 447; mnioides, 486; pallidus, 486, 505; paradoxus, 486, 505
Texas. a Trichostomum new to the United States, 80; macrolichens of Big Bend National Park, 296; mosses of Big Bend National Park, 269; new and additional lichen records from, 224
Thamnium andinum, *151*, *169*; fasciculatum, *151*, *169*; lombrophyllaceum, *151*, *169*; longirostre, *151*, *169*; ramosissimum, *151*, *169*; thyrsoides, *151*, *169*
Thamnolia subuliformis, 455; vermicularis, 55-, 455
thamnolic acid, 385-
Thchad. *See* Africa
Thelia asprella, 95-
thiophanic acid, 385-
Thomas, P. E. & P. L. Redfearn. Bryophytes of the Interior Highlands XVI. New Distributional Records for Arkansas and Missouri, 100
Thomson, J. W. & T. H. Nash III. Three New Lichens from the Southwest: Xanthoria concinna sp. nov., Lecanora collatolica sp. nov. and Toninia conglomerata, 350
Thuidiaceae, 154, 292
Thuidium abietinum, 447; brasiliense, 154; breviacuminatum, 154; ciliatum, 154; delicatulum, 1-, 95-, *154*, *169*, 222-; delicatulum var. radicans, 447; firmulum, *154*, *169*; involvens, 154, 511; latopulvinatum, 154; leptocladum, 154; minutulum, 154, 169; ochraceum, 154; pauperum, *154*, *169*; peruvianum, 154, 168; pusillum, 154; recognitum, 447; recognitum var. delicatulum, 154, 169; scabrosulum, 154; schistocalyx, 154; tamariscinum, 86; tomentosum, 511; yungarum, 154
Thyrea nigritella, 312; pulvinata, 312
Tilia, 241-; americana, 241-; cordata, 86-
Timmia austriaca, 447, 486; megapolitana var. bavarica, 447

Timmiella anomala, 516
Toivonen, S. *See* Valanne N., et al.
Tomenthypnum nitens, 447, 486
Toninia coeruleonigricans, 172-, 455; conglomerata, 353*
Tortella arctica, 208-, 447, 486, 505; caespitosa, *135*, *169*; contortifolia, 135, 169; fragilis, 321-, 447; germainii, 135, 161; humilis, 135, 169, 284; inclinata, 447; mollissima, 512; pilcomayica, 135; pileomayica, *135*; sp., 55-; tortuosa, 447, 505
Tortula aculeata, 135; andicola, 136, 161; angustifolia, 136, 162; armata, *134*, *169*; atrovirens, 516; aurea, 285; bartramii, 217, 285; bipedicellata, 136; brunnea, 136, 161; buchtienii, 136; campylocarpa, *133*, *169*; ciliata, 136; contortifolia, *135*, *169*; denticulata, 136; fragilis, 136, 285; glacialis, 136; goudotii var. boliviana, 136; lacerifolia, 136; latifolia, 86-; leiostoma, 136; ligulata, 136; limbata, *133*, *169*; linguifolia, 136; minima, 136; mniadelphus, 136, 161; mniifolia, 136; mucronifolia, 321, 447, 486, 505; norvegica, 447, 505; obtusissima, 285; pagorum, 217, 285; papillosa, 217, 285; percarnosa, 136; pichinchensis, 136, 169; polylepidis, 136; polyseta, 136, 161; pulvinata, 86-; rhizzophylla, 136, 227-; robusta var. runcinata, 136, 169; runcinata, *136*, *169*; ruraliformis, 217; ruralis, *169*, 208-, 285, 339-, 447, 486; ruralis var. spiralis, 136, 169; scabrinervis, 136; semivaginata, *134*, *169*; serripungens, 136; serripungens var. exesa, 136; sordida, 136; sp., 5-; spiralis, *136*, *169*; subulata, 86-; vectensis, 227-; viridula, 136, 161; xerophila, 136
Trebouxia, 350-
Trematodon bolivianus, 131; brevicollis, 447, 486, 505; reflexus, 131
Trichocolea tomentella, 1-
Trichopilum polycarpum, *160*, *169*
Trichosteleum ambiguum, 158; arrectum, 158; fluviale, 159; microcarpum, 159
Trichostomopsis australasiae, 136, 285; australis, 217; brevifolia, 269-; diaphanobasis, 269-
Trichostomum aeneum, 229; apophysatulum, 136; brachydontium, 80-, 285; campylocarpum, *133*, *169*; campylopyxis, *133*, *169*; canaliculatum, *136*, *169*; challaense, *134*, *169*; chilensis, *133*, *169*; compactulum, 136; crispulum var. cucullatum, 81*, 82; cucullatum, 82; cuspidatissimum, 505; edentulum, 136; fallax, *135*, *169*; ferrugineum, *134*, *169*; gracilescens, *134*, *169*; grimmioides, *135*, *169*; involutum, 136, 169; jamaicense, 285; mandonii, *169*; paludicola, 136, 161; pomangium, 136; quitense, *136*, *169*; quitense var. longifolium, *136*, *169*; rigidiusculum, 136; semimarginatum, *134*, *169*; syrrhopodontoides, *135*, *169*; tophaceum, 134, 169
Tridontium tasmanicum, 16-
Trifolium spp., 172-
Trillium grandiflorum, 1-
Triquetrella cucullata, *135*, *169*
Tristichium lorentzii, 128; lorentzii var. apodum, 128; mirabile, 128
Tritomaria exsecta, 184, 427; exsectiformis, 184, 427; polita, 184, 427; quinquedentata, 184, 427; scitula, 184, 427
Trochobryum, 231-
Tsuga canadensis, 385-; caroliniana, 222-

Udar, R. & D. K. Singh. A New Cyathodium from India, 234
Ulex europaeus, 172-
Ulmus, 241-; americana, 64-, 241-; glabra, 86-
Ulota curvifolia, 321-, 506
Ulothrix, 22-
ultrastructure. antheridia of Polytrichum, 466; chloroplasts in germinating spores of Polytrichum, 332; germinating spores of Dicranum, 188; leaf sections and protonemata of mosses, 339; plastids in Marchantia, 22
Ulva californica, 397; tesselata, *396*; vextata, *397*
Umbilicaria cylindrica, 455; deusta, 455; havaasii, 455; hyperborea, 455; proboscidea, 455; vellea, 312
Urceolaria mutabilis, *399*
uses of bryophytes, 95
Usnea arizonica, 312; cavernosa, 103-; cirrosa, 312; comosa, 172-, *312*; diplotypus, 312; florida, 172-; fulvoreagens, 402; glabrata, 455; glabrescens, 455; glabrescens var. fulvoreagens, *402*; hirta, 312; mutabilis, 1-; sorediifera, *402*; spp., 172-; subfloridana, 312; subfusca, 296-; sulcata, 360; trichodea, 402
usnic acid, 224-, 296-, 364-, 385-
Utah. Pterygoneurum lamellatum in, 221

Vaccinium palustris, 47-; spp., 222-; uliginosum, 47-
Valanne, N., S. Toivonen & R. Saarinen. Ultrastructural Changes in Germinating Dicranum scoparium: A Moss Containing Protein Storage Material, 188
Valanne, N. *See also* Karunen, P. et al.
vegetative reproduction. in Ramalina subleptocarpha, 364
Venezuela. distribution of Rozea in, 413
Verrucaria maura, 402
Verrucidens, 231-

Vesicularia amphibola, 160, 163; poeppigiana, 160, 169; vesicularis, 160, 163, 511; vesicularis var. poeppigiana, *160, 169*
Virginia. Hypotrachyna showmanii in, 80
Vitt, D. H. Mosses New to the Yukon from the Ogilvie Mountains, 501
Vitt, D. H. & W. B. Schofield. Seligeria careyana, a New Species from the Queen Charlotte Islands, Western Canada, 231
Vitt, D. H. *See also* Douglas, G. W. & D. H. Vitt
Vitt, D. H. *See also* Hong, W. S. & D. H. Vitt
Voitia hyperborea, 486

wall development in spermatogenous cells of Polytrichum, 466
Washington. distribution of Ramalina subleptocarpha in, 367*, 368; Schofieldia, a new hepatic in, 314; Sphagnum in, 508
Weber, D. P. *See* Brassard, G. R., D. P. Weber & A. J. Fife
Webera apolensis, *141, 169*; clavicaulis, *141, 169*; cruda, *141, 169*; loriformis, *141, 169*; papillosa, *141, 169*; spectabilis, *142, 169*; subleptopoda, *142, 169*
Weisia controversa, 1-, 86, 106-, 136, 169, 286, 447; glauca, 217, 286; ligulifolia, 286; longidentata, *136, 169*; rostellata var. phascoides, 384; spp., 80-; tortivelata, 136; viridula, *136, 169*
Weissia. *See* Weisia
West Indies. Notes on Pottiaceae, 227-
Wetmore, C. M. Macrolichens of Big Bend National Park, Texas, 296
Wilkinson, R. E. *See* Karunen, P. et al.
Willia, 122-
Williamsia tricolor, *20, 134, 169*
Williamsiella tricolor, *20, 134, 169*
Wilsoniella flaccida, 131, 169
Wistendahl, W. A. *See* McFarland, K. D. & W. A. Wistendahl
Wollnya stellata, *140, 169*
xanthone, 385-
Xanthoparmelia chlorochroa, *401*; dissensa, 224-; psoromifera, 224; **nigropsoromifera**, comb. nov., 225; weberi, 225
Xanthoria candelaria, 313, 360, 403, 455; **concinna**, sp. nov., 350, 351*; elegans, 456; fallax, 313; parietina, 86-; polycarpa, 313, 350-, 456
Xylographa abietina var. abietina, 403; parallela, *403*; spilomatica, *403*; vitiligo, 403

Yucca spp., 269-
Yukon. *See* Canada

Zander, R. H. Notes on Pottiaceae in Middle America, 227
Zander, R. H. & E. Hegewald. Leptodontiella, gen. nov. and Leptodontium from Peru, 16
zeorin, 296-, 364-
Zygodon andinus, *147, 169*; basidentatus, *147, 169*; brevipes, 146, 169; caldensis, *147, 169*; coraniensis, *146, 169*; denticulatus, *147, 169*; fasciculatus, 146; ferrugineus, *147, 169*; ferrugineus var. viridis, *147, 169*; goudotii, 146, 169; gracillimus, 146, 169; inconspicuus, *146, 169*; intermedius, 146, 169; liliputanus, *146, 169*; linguiformis, *146, 169*; macrophyllus, 146; mandonianus, *133, 169*; obtusifolius, 146, 169; oeneus, 146; ovalis, *146, 169*; paucidens, *147, 169*; perichaetialis, 146; peruvianus, 147, 169; pichinchensis, 147; pilosulus, 147, 169; pyriformis, 147; ramulosus, 147; recurvifolius, *147, 169*; reinwardtii, 147, 169; reinwardtii var. subintegrifolius, 147, 169; rigens, *147, 169*; stenocarpus, *147, 169*; subdenticulatus, *147, 169*; *subrecurvifolius*, 147, 169, var. recurvatus, *147, 169*; subrecurvifolius var. viridis, 147, 169; vestitus, 147; viridissimus, 86-, 215-